

MAPK inhibitors induce serine peptidase inhibitor Kazal type 1 (SPINK1) secretion in BRAF V600E mutant colorectal adenocarcinoma

Kati Räsänen^{1#}, Kien X. Dang², Harri Mustonen³, Tho H. Ho⁴, Susanna Lintula¹, Hannu Koistinen¹, Ulf-Håkan Stenman¹, Caj Haglund^{3,5,6*} and Jakob Stenman^{2,7*}

¹Department of Clinical Chemistry, Medicum, University of Helsinki and Helsinki University Hospital, 00014 Helsinki, Finland

²Minerva Foundation Institute for Medical Research, Helsinki, 00290, Finland

³Department of Surgery, University of Helsinki and Helsinki University Hospital, Helsinki, 00029 HUS, Finland

⁴Department of Genomics, BPARC, Vietnam Military Medical University, Hanoi, 10000 Hanoi, Vietnam

⁵Research Program Unit, Translational Cancer Biology, University of Helsinki, 00014 Helsinki, Finland

⁶Department of Pathology, University of Helsinki and Helsinki University Hospital, 00014 Helsinki, Finland

⁷Department of Women's and Children's Health, Karolinska Institutet, 17176 Stockholm, Sweden

*Equal contribution

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/1878-0261.12160

Molecular Oncology (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

#Corresponding author

Kati Räsänen, PhD

Department of Clinical Chemistry, Medicum

University of Helsinki

00014 Helsinki

Finland

Email kati.rasanen@helsinki.fi

Tel. +358 947171725

Keywords: BRAF V600E, SPINK1, colorectal cancer, biomarker, inhibitor, trametinib

Running title: SPINK1 secretion in BRAF V600E colon cancer

Abbreviations: ADAM, A Disintegrin and Metalloproteinases; ATF-4, activating transcription factor 4; CRC, colorectal cancer; EGFR, epidermal growth factor receptor; ExBP-RT, Extendable Blocking Probe Reverse Transcriptase; FFPE, formalin-fixed paraffin embedded; IFMA, immunofluorometric assay; IHC, immunohistochemistry; IQR, inter quartile range; MAPK, mitogen-activated protein kinase; qPCR, quantitative PCR; SPINK1, serine peptidase inhibitor Kazal type 1; TIMP1, tissue inhibitor of metalloproteinases 1

Abstract

The mitogen-activated protein kinase (MAPK) pathway plays a central role in colorectal cancers (CRC). In particular, BRAF V600E-mutant tumors, which represent around 10% of CRCs, are refractory to current therapies. Over-expression and secretion of serine peptidase inhibitor Kazal type

1 (SPINK1) is observed in around 50% of CRCs and its serum level can be used as a biomarker for poor prognosis. Utilizing a recently developed Extendable Blocking Probe assay, we analyzed the BRAF mutation status in a CRC patient cohort (N=571) using tissue-derived RNA as the starting material. From the same RNA samples we measured the relative SPINK1 expression levels using a quantitative real-time PCR method. Expression of mutant BRAF V600E correlated with poor prognosis, as did low expression of SPINK1 mRNA. Further, BRAF V600E correlated negatively with SPINK1 levels. In order to investigate the effect of MAPK pathway targeted therapies on SPINK1 secretion, we conducted *in vitro* studies using both wild-type and V600E colorectal cancer cell lines. BRAF inhibitor vemurafenib, and subsequent MAPK pathway inhibitors trametinib and SCH772984, significantly increased SPINK1 secretion in V600E CRC cell lines Colo205 and HT-29 with concomitant decrease in trypsin-1 and -2 secretion. Notably no SPINK1 increase or trypsin-1 decrease was observed in BRAF wild-type CRC cell line Caco-2 in response to MAPK pathway inhibitors. In further mechanistic studies, we observed that only trametinib was able to diminish completely both MEK and ERK phosphorylation in the V600E CRC cells. Furthermore, the key regulator of integrated stress response, activating transcription factor 4 (ATF-4), was downregulated both at mRNA and protein level in response to trametinib treatment. In conclusion, these data suggest that sustained inhibition of not only MAPK pathway activation, but also ATF-4 and trypsin, might be beneficial in therapy of BRAF V600E-mutant CRC and that SPINK1 levels may serve as an indicator of therapy response.

1. Introduction

Colorectal cancer (CRC) is the third most common cancer in men and the second in women worldwide. While about 20% of patients have disseminated disease at diagnosis, part of the patients initially classified as having regional disease and even some of those with local disease will develop a recurrence and eventually die from cancer. Overall 5-year survival is 50-60% (Siegel et al., 2012). Current FDA-approved molecularly targeted therapies for metastasized CRC include several

monoclonal antibodies against vascular endothelial growth factor and epidermal growth factor receptor (EGFR), a multi-kinase small molecule inhibitor regorafenib and a nucleoside analog trifluridine/tipiracil (Moriarty et al., 2016). Currently available EGFR antibodies cetuximab and panitumumab bind to the extracellular domain of EGFR resulting in receptor internalization and blockage of signaling. Mutations in the RAS family of proto-oncogenes (KRAS, NRAS, HRAS) result in constitutive activation of the mitogen-activated protein kinase (MAPK) pathway signaling independent of activation of receptor tyrosine kinases such as EGFR. Thereby, mutations in KRAS or NRAS cause intrinsic resistance to EGFR-targeted therapies (Semrad and Kim 2016).

V600E mutation in BRAF, which is immediately downstream of RAS, has also been proposed to cause resistance to anti-EGFR antibodies. This mutation is found in about 10% of CRCs leading to an aggressive subtype, for which there are no effective oncological therapy (Dienstmann and Tebernero 2016). Unexpectedly specific BRAF V600E inhibitors, like vemurafenib that is highly effective in melanoma, do not benefit CRC patients. Subsequently it was shown that intrinsic resistance to vemurafenib in CRC is caused by EGFR auto-activation through an unknown mechanism (Prahallad et al., 2012). In recent clinical trials combination of a BRAF V600E inhibitor with a MEK (Corcoran et al., 2015) or a PI3K (Elez et al., 2015) inhibitor showed clinical benefit. Therefore, sustained MAPK inhibition appears to be a critical determinant of the clinical benefit and, differing from melanoma, it seems that only combinations are able to generate therapeutic effects in CRC (Dienstmann and Tebernero 2016).

Over-expression and secretion of serine peptidase inhibitor Kazal type 1 (SPINK1, also known as tumor-associated trypsin inhibitor, TATI or pancreatic secretory trypsin inhibitor, PSTI) is observed in a variety of cancers (Räsänen et al., 2016a). In CRC, increased SPINK1 serum levels are found in around 50% of patients and it is an independent prognostic factor (Gaber et al., 2010). However, the prognostic value of tissue expression of SPINK1 in colorectal cancer is controversial, as depending on study it has been predictive of either poor or good prognosis, or non-significant (Chen et al., 2015, Koskensalo et al., 2013, (Ida et al., 2015)). We have previously shown that concomitant SPINK1 and EGFR expression in CRC tissue predicts favorable prognosis (Koskensalo et al., 2013);

Molecular Oncology (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd.

and Koskensalo and colleagues (Koskensalo et al., 2012) and Chen and colleagues (Chen et al., 2016) showed that high tissue expression of SPINK1 in CRC correlated with a better prognosis.

In addition to intrinsic resistance, acquired resistance presents a clinical problem as a majority of the patients who are treated with molecularly targeted treatments relapse within a year. Therefore, both novel methods and novel biomarkers that predict treatment-response are needed for stratification of patients in order to select appropriate therapy. In the current study, we investigated the expression of BRAF V600E mutations in a CRC cohort of 571 patients using a novel Extendable Blocking Probe Reverse Transcriptase (ExBP-RT) assay that we recently developed. ExBP-RT is an ultra-high selective method and allows for analysis of expressed mutations at the RNA level (Ho et al., 2015). This method therefore enables analysis of mutational status irrespective whether the mutation is inherited or acquired. The use of tumor tissue RNA instead of DNA as the starting material enabled us to correlate the mRNA expression of BRAF V600E mutations directly to SPINK1 mRNA expression level analyzed by qPCR from the same tumor samples. Further, using BRAF wild type and V600E colorectal cancer cell lines, we studied the effects of MAPK inhibitors on SPINK1 secretion *in vitro*.

2. Materials and methods

2.1 RNA samples

RNA was extracted as described (Ho et al., 2015) from formalin-fixed paraffin embedded (FFPE) samples from patients who were operated for histologically confirmed colorectal cancer at the Department of Surgery, Meilahti Hospital, Helsinki University Hospital between 1987-2003. In total 571 patients' samples were available for this study. The use of clinical samples for this purpose was approved by the Surgical Ethics Committee of Helsinki University Hospital and the National Supervisory Authority of Welfare and Health and collected from the archives of the Department of Pathology, Helsinki University Hospital. All RNA samples were quantified with a NanoVue spectrophotometer (GE Healthcare, Waukesha, WI). Table 1 describes the clinicopathological features of the cohort.

Molecular Oncology (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd.

2.2 ExBP-RT assay

The Extendable Blocking Probe method (Ho et al., 2015) was employed for ultrasensitive detection of the BRAF V600E gene mutation. Using RNA templates, this novel method allows for detection of expressed mutations in at least a 1000 times higher background of the corresponding wild type alleles. The principles of ExBP-RT assays and reaction set-up procedures for multiplex detection of BRAF V600E mutation were as described in the original paper (Ho et al., 2015).

RNA extracted from FFPE samples was diluted to 100 ng/ μ l in DEPC H₂O for the ExBP-RT assay, before the allele-specific reverse transcription reaction. RNA extracted from Colo205 (BRAF V600E mutant) and A549 (BRAF wild-type) cell lines were used as positive and negative controls, respectively, in ExBP-RT assays of BRAF mutation detection. All control RNAs were extracted from cultures cells using RNA/DNA purification Kit (Norgen Biotek, Thorold, Canada), quantified with a NanoVue spectrophotometer (GE Healthcare) and diluted to 100 ng/ μ l in DEPC H₂O.

Using cDNA products of the ExBP-RT assays as template, the real-time PCR amplification was performed to detect/quantify the expression of mutant BRAF V600E. QuantiTect Probe PCR Kits (Qiagen, Hilden, Germany) were used for these probe-based real-time PCR assays according to the manufacturer's instructions in a 10 μ L reaction volume. A common reverse primer was designed to target the 5'-prime tail of all mutation-specific ExBP-RT products. The expression levels of total BRAF genes (including V600E mutant and its wild-type segments) were also determined in each sample for normalization using QuantiTect SYBR Green PCR Kits (Qiagen) according to the manufacturer's instructions in a 10 μ L volume. The sequences and concentration of qPCR primers and probes are provided in Table 2. The same thermocycling conditions were used for both probe-based and SYBR Green-based real time quantitative PCR (qPCR): 95°C for 15 min, 45 cycles at 94°C for 10 sec, at 60°C for 45 sec. Following SYBR Green-based qPCR, the specificity of the amplification products was verified by melting curve analysis. All qPCR assays were run on a LightCycler 480 II Real-Time PCR Instrument (Roche Applied Science, Mannheim, Germany) with a 384-well white-plate (Roche Applied Science). All mutation, wild type and H₂O controls of each experiment were checked to verify the results in both ExBP-RT and qPCR assays. Threshold cycle (Ct) values of qPCR

Molecular Oncology (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd.

were calculated automatically using the Absolute Quantification Analysis with the Fit Points Method, which is built in the LightCycler 480 II system. The method allows to set the noise-band and the threshold line in order to discard uninformative background noise.

2.3 Real-time quantitative PCR

For FFPE samples 500 ng of total RNA was reverse transcribed with 100 U Revert Aid Premium Reverse Transcriptase (Thermo Fisher Scientific, Waltham, MA) using 4 pmol of gene-specific antisense primers for *SPINK1* and *RPL13A* (see below for sequences), 0.5 mM dNTP mix and 20 U Ribolock RNase Inhibitor (all from Thermo Fisher Scientific). Possible contamination of RNA in FFPE-extracted samples with *SPINK1* or *RPL13A* DNA was excluded by subjecting each sample to RT reaction without Revert Aid Premium Reverse Transcriptase. Real-time qPCR was performed with a LightCycler 480 II instrument using a 384-well thermal block (Roche Applied Science) with SensiFAST SYBR No-ROX Kit (Bioline, London, UK). *SPINK1*, *PRSS1* and *PRSS2* qPCR from cell lines was performed using conditions described previously (Räsänen et al., 2016b). The following primers, purchased from TAG Copenhagen (Copenhagen, Denmark) and verified earlier (Räsänen et al., 2016b), were used: *SPINK1* forward 5'- TGT CTG TGG GAC TGA TGG AA, *SPINK1* reverse 5'- GCC CAG ATT TTT GAA TGA GG, *PRSS1* forward 5'-CCA CCC CCA ATA CGA CAG GAA G, *PRSS1* reverse 5'-GCG CCA GAG CTC GCA GT, *PRSS2* forward 5'-CCA AAT ACA ACA GCC GG, *PRSS2* reverse 5'-AGT CGG CAC CAG AAC TCA GA, *RPL13A* forward 5'-AGA TGG CGG AGG TGC AG and *RPL13A* reverse 5'- GGC CCA GCA GTA CCT GTT TA.

Following SYBR Green-based qPCR, the specificity of the amplification products was verified by melting curve analysis and a control sample was included in every run to confirm inter-assay reproducibility. All reactions were run in duplicate and for all samples RT- controls were run to exclude possible DNA contamination. Relative expression of target gene mRNA referenced to *RPL13A* housekeeping gene was calculated using the $\Delta\Delta C_t$ method (Livak et al., 2001).

Human Transcription Factors (Non-HOX) TaqMan Array (Thermo Fisher Scientific) that contains 92 assays to non-HOX transcription factor -associated genes and 4 assays to endogenous control genes was performed according to manufacturer's protocol.

2.4 Immunohistochemistry

SPINK1 immunohistochemistry (IHC) was performed as described recently (Räsänen et al., 2016b). Briefly, FFPE sections were deparaffinized with xylene and rehydrated in ethanol series. Mouse monoclonal anti-SPINK1 (2 µg/ml, in-house clone 6E8) (Osman et al., 1993) was incubated overnight at 4 °C. Isotype control antibody was Mouse IgG #I-2000 (Vector Laboratories, Burlingame, CA). Mach 4 Universal AP-Polymer kit (Biocare Medical, Concord, CA) was used for detection, and the visualization signal was developed with Vector Red Alkaline Phosphatase Substrate kit (Vector Laboratories).

2.5 Cell culture

BRAF V600E colorectal cancer cell lines Colo205 and HT-29 (American Type Culture Collection, Manassas, VA), and BRAF wild type cell lines Caco-2, and SW-480 (Sigma-Aldrich, St. Louis, MO), were cultured and authenticated as previously described (Räsänen et al., 2016b). Melanoma cell lines SK-MEL-2 (BRAF wild type) and SK-MEL-5 (BRAF V600E) were purchased from American Type Culture Collection and grown in EMEM. All cell lines were cultured at +37 °C in 5% CO₂ atmosphere and supplemented with 5% fetal bovine serum (Biowest, Nuaille, France), 0.3 mg/ml glutamine, 100 µg/ml streptomycin, and 100 U/ml penicillin (all from Lonza, Basel, Switzerland). Cells were used until passage number 20 and routinely tested for mycoplasma.

2.6 Inhibitors

The following inhibitors were purchased from Selleck Chemicals (Munich, Germany) and dissolved in DMSO according to manufacturer's instructions: gefitinib (EGFR), GW5074 (CRAF), LY294002 (AKT), PD98059 (MEK1), salisarib (RAS), SCH772984 (ERK1/2), trametinib (MEK1/2), vemurafenib (BRAF) and zoledronic acid (RAS and RHO).

2.7 Immunofluorometric assay

Time-resolved immunofluorometric assays (IFMA) developed in-house for SPINK1, trypsinogen-1 and trypsinogen-2 (Janeiro et al., 2012, Paju et al., 2001, Itkonen et al., 1990) were performed as described previously (Räsänen et al., 2016b). The concentrations of secreted proteins were measured from 24, 48 and 72h conditioned media with or without various inhibitor treatments. The detection ranges for the IFMAs are the following: SPINK1

0.5 – 90 ng/ml, trypsinogen-1 1.6 – 400 ng/ml and trypsinogen-2 2 – 500 ng/ml.

2.8 Western blotting

Samples for Western blot analysis were harvested as described (Räsänen et al., 2008). Samples were run on 4-12% gradient gels (Life Technologies, Carlsbad, CA). Using Trans-Blot Turbo system, proteins were transferred to nitrocellulose membrane (both from Bio-Rad, Hercules, CA) and blocked with 5% (w/v) non-fat powdered milk in TBS (20 mM Tris-HCl pH 7.5, 150 mM NaCl and 0.1% Tween-20). Immunoreactive proteins were visualized with appropriate primary and secondary antibodies using ECL detection (Bio-Rad).

The following primary antibodies were used at manufacturers' recommended dilutions: rabbit monoclonal anti-phospho-p44/42 MAPK (ERK1/2, Thr202/Tyr204), rabbit polyclonal anti-p44/42 MAPK (ERK1/2), rabbit monoclonal anti-phospho-MEK1/2 (Ser217/221), mouse monoclonal anti-

Molecular Oncology (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd.

MEK1/2 (L38C12), rabbit polyclonal anti-phospho-STAT3 (Y705), mouse monoclonal anti-STAT3 (124H6) and rabbit monoclonal anti-ATF4 (D4B8) (all from Cell Signaling Technology, Danvers, MA). Rabbit polyclonal anti-GAPDH was from Sigma-Aldrich. The secondary antibodies used in Western blotting were affinity-purified horseradish peroxidase coupled anti-rabbit IgG H+L and anti-mouse IgG H+L (both from Jackson ImmunoResearch, Suffolk, UK).

2.9 Statistical analysis

Results are given as number of patients and percentage or mean and SD or median and range or interquartile range (IQR). The Fisher's exact-test and the linear-by-linear association test were used to assess associations between clinicopathological variables and mutation status or mRNA expression. The relative SPINK1 mRNA expression was dichotomized at 87.5% percentile. Differences in continuous variables between different groups were tested with the unpaired T-test or with the Mann-Whitney test or in the case of ordinal grouping variable with the Jonckheere-Terpstra test. Survival analysis was performed with the Kaplan-Meier method, and the log rank test was used to compare the groups. The Cox regression proportional hazard model was used for uni- and multivariate survival analysis. Multivariate analysis was adjusted for age, gender, stage and location. Interaction terms were considered. The Cox model assumption of constant hazard ratios over time was tested. A time-dependent covariate was included separately for each testable variable at a time. A time-dependent correction factor was included into the models, if the hazard ratio was not constant over time. The Spearman's rho correlation coefficient was calculated to assess correlation between continuous and ordinal variables. All *in vitro* experiments were done in duplicate and repeated three times. P-values of less than 0.05 were considered to be statistically significant and two tailed tests were used. Statistical analyses were carried out with SPSS (version 24, IBM, NY) and GraphPad Software (La Jolla, CA).

3. Results

3.1 BRAF V600E mutation detection by ExBP-RT assay

BRAF V600E mutation is known to be a strong marker of poor prognosis in metastatic CRC. The prevalence of this mutation is around 10% (Dienstmann and Tebernero 2016). We detected expressed BRAF V600E mutations in 8% of the samples in our cohort, and presence of BRAF V600E correlated significantly with poor prognosis (the Log Rank test, $p < 0.001$). Patients with no detected expression of V600E mutations had mean survival time of 16.1 (95% CI, 15.0-17.2) years compared to 11.1 (8.0-14.2) years for patients with expressed mutations (Figure 1). In univariate Cox analysis the hazard ratio for expressed BRAF V600E mutations was 2.1 (95% CI, 1.5-3.0, $p < 0.001$) and in multivariate model after adjusting for age, gender, stage and location it was 2.8 (95% CI, 1.8-4.4, $p < 0.001$) (Table 3).

3.2 Low expression of tumor SPINK1 mRNA associates with poor prognosis

First we correlated the SPINK1 mRNA qPCR data to the previously published (Koskensalo et al., 2012) immunohistochemistry result of this cohort. The relative expression level of SPINK1 measured by qPCR significantly correlated with the previous IHC results (Spearman's rho 0.366, $p < 0.001$, $N=242$). Representative images of SPINK1 IHC and corresponding relative SPINK1 mRNA levels are shown in Figure 2A. Further, in line with the reported result of the IHC staining where low SPINK1 immunoreactivity was an independent prognostic factor for adverse outcome (Koskensalo et al., 2012), low SPINK1 mRNA expression was associated with poor prognosis (Figure 2B, the Log Rank test, $p=0.033$). This was more prominent in patients having disease on left side (Figure 2C, the Log Rank test $p=0.004$). Patients with high SPINK1 mRNA level (>2.4 on relative scale) and left side disease had mean survival time of 17.4 (95% CI, 14.9-19.8) years compared to 14.5 (95% CI, 12.9-16.1) years for patients with lower SPINK1 mRNA level (≤ 2.4). In univariate Cox analysis the hazard ratio for SPINK1 mRNA level was 0.51 (95% CI, 0.29-0.91, $p=0.023$, relative mRNA

expression =>2.4 vs <0.5), and 0.43 (95% CI, 0.22-0.84 p=0.014) in multivariate model adjusted for age, gender, stage and location (Table 3).

3.3. Correlation of expressed BRAF V600E mutations to SPINK1 expression

As the independent analyses of expressed BRAF V600E mutations and low SPINK1 expression were indicative of poor prognosis, we analyzed the correlation between these two biomarkers. Expressed BRAF V600E mutation correlated negatively with both SPINK1 mRNA expression level (Spearman's rho -0.19, p<0.001) and with the previously published (Koskensalo et al., 2012) IHC results (Spearman's rho -0.21, p<0.001). The relative SPINK1 mRNA expression was lower in samples with expression of mutated BRAF V600E (median 0.4, IQR 0.1-0.6) than in samples with BRAF wild type expression only (median 0.8, IQR 0.3-1.5, p=<0.0001, Mann-Whitney test). The hazard ratio of high SPINK1 mRNA level (>2.4) in patients with expression of wild type BRAF to patients with expressed BRAF V600E mutations was 0.09 (95% CI, 0.03-0.27) after adjusting for age, gender, stage and location. All patients, except one, with expressed BRAF V600E mutations, had low SPINK1 mRNA levels (<=2.4 relative expression).

3.4 Effect of MAPK inhibitors on SPINK1 secretion

Next we analyzed the levels of secreted SPINK1 and its putative serine protease targets trypsin-1 and trypsin-2 in a panel of colorectal cancer and melanoma cell lines harboring either wild type or V600E BRAF. Table 4 shows the basal levels of SPINK1, trypsin-1 and trypsin-2 at 72h time point in a panel of cell lines and their respective BRAF status. Notably, neither of the melanoma cell lines secreted SPINK1 or trypsins.

In order to study the effects of various MAPK pathway inhibitors on the SPINK1 levels, colorectal cancer cell lines were treated with the following compounds: gefitinib, GW5074, LY294002, PD98059, salisarib, SCH772984, trametinib, vemurafenib and zoledronic acid. In BRAF Molecular Oncology (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd.

V600E colorectal cancer cells lines Colo205 and HT-29, inhibitors affecting the MAPK pathway at or below BRAF resulted in over 2-fold dose-dependent increase in SPINK1 secretion measured at 72h time point (Figure 3 A and B). This effect was seen with the BRAF inhibitor vemurafenib and subsequent MAPK pathway inhibitors trametinib (MEK1/2 inhibitor) and SCH772984 (ERK1/2 inhibitor), but interestingly not with the CRAF inhibitor GW5074 or with PD98059 that is a non-ATP competing MEK antagonist specifically inhibiting MEK1-mediated activation of the MAPK pathway. Further, inhibitors upstream of RAF, such as the RAS inhibitor salisarib, RAS/RHO inhibitor zoledronic acid or EGFR inhibitor gefitinib did not induce SPINK1 secretion in BRAF V600E CRC cells. These effects were not seen in the BRAF wild type CRC cell line Caco-2 (Figure 3 C). Akt inhibitor LY294002 did not affect SPINK1 levels in any of the tested cell lines.

3.5 MAPK inhibitors induce SPINK1 and concomitantly downregulate trypsin-1 and -2 in BRAF V600E cells

To further elucidate the effects of the MAPK pathway inhibitors on CRC cells, using the minimum dose that induced SPINK1 secretion in Colo205 in HT-29 cells (60 nM), we measured the levels of SPINK1 in a time-dependent manner. At 48h and 72h time points in both cell lines vemurafenib, trametinib and SCH772984 resulted in a statistically significant increase in SPINK1 secretion ($p < 0.05$) compared to control, as measured by IFMA (Figure 4 A and B, top panels). Corroborating the immunoassay results, SPINK1 mRNA levels were increased at the 72h time point in response to vemurafenib, trametinib and SCH772984 in both cell lines (Supplemental Figure 1 A).

Furthermore, as SPINK1 is a putative trypsin inhibitor, we investigated whether the MAPK inhibitors affected endogenous trypsin levels in the CRC cell lines. In the BRAF V600E Colo205 cells vemurafenib, trametinib and SCH772984 led to a statistically significant ($p < 0.05$) decrease in trypsin-1 (Figure 4 A, mid panel) and trypsin-2 (Figure 4 A, bottom panel) levels at 72h time point, suggesting inverse regulation between SPINK1 and its target proteases. In the other BRAF V600E cell line HT-29 (Figure 4 B) vemurafenib and SCH772984 led to statistically significant ($p < 0.05$)

decrease in trypsin-1 and -2 levels, whereas in trametinib-treated cells trypsin-1 and -2 levels did not decrease. These data were confirmed by qPCR analyses of the mRNA levels of PRSS1 (trypsin-1) and PRSS2 (trypsin-2) (Supplemental Figure 1 B and C).

In the BRAF wild type cell line Caco-2, level of SPINK1 was slightly increased in response to trametinib at 48h time point, but not in response to vemurafenib and SCH772984, as analyzed by IFMA (Figure 4 C). Trypsin-1 levels were not affected by inhibitors (Figure 4 C, right panel) and trypsin-2 was not detected in the Caco-2 cell conditioned media by IFMA. The results were confirmed by qPCR analyses of the mRNA levels of SPINK1, PRSS1 (trypsin-1) and PRSS2 (trypsin-2) indicating correlation between mRNA expression and secretion of these proteins (Supplemental Figure 1).

3.6 Trametinib diminishes ERK1/2 and MEK1/2 phosphorylation and downregulates ATF-4

To elucidate the signaling events that led to increased SPINK1 expression in response to MAPK inhibitors, we studied the phosphorylation status of MEK1/2 and ERK1/2 at 24h time point (Figure 5). In both Colo205 and HT-29 cells trametinib diminished completely the phosphorylated forms of ERK1/2 (Thr202/Tyr204) and MEK1/2 (Ser217/221). However, vemurafenib and SCH772984 reduced phosphorylation of ERK1/2 and MEK1/2 to a much lesser extent compared to trametinib. These results were in line with the observed increase in SPINK1 levels in these cells. Further, PD98059, which did not affect SPINK1 expression (Figure 4) did not affect the phosphorylation status of ERK1/2 or MEK1/2 in either CRC cell line.

Next we performed a TaqMan non-HOX transcription factor array in order to identify which transcription factor(s) is responsible for the increased SPINK1 expression. As trametinib caused the biggest increase in SPINK1 level in both cell lines, we used it at 60 nM concentration and harvested RNA at 24h time point. Interestingly, none of the transcription factors included in the array were significantly induced in the trametinib-treated Colo205 and HT-29 cells (Figure 6A). The only

transcription factor with a change in its mRNA level in the trametinib-treated sample compared to the DMSO control was activating transcription factor 4 (ATF-4), a transcription factor linked to integrated stress response (IRS) (Pakos-Zebrucka et al., 2016). The decrease in ATF-4 caused by trametinib was further confirmed by Western blotting (Figure 6B).

As we have previously shown that interleukin-6 induces SPINK1 expression in Colo205 and HT-29 cells via STAT3 phosphorylation (Räsänen et al. 2016b), we investigated whether this was also the case with the MAPK inhibitors. Phosphorylation of STAT3 was not induced in response to the inhibitors (data not shown) and the total STAT3 levels remained constant (Supplementary Figure 2).

4. Discussion

CRC patients with the BRAF V600E mutation present a clinical challenge, as no effective treatments have been found for this subpopulation. In keeping with previously published results (Barras 2015), patients with this mutation have a decreased survival rate in our cohort. Several publications have shown that SPINK1 plays a role in the tumorigenesis of CRC, in particular at the later stages (Räsänen et al., 2016a). In this study we demonstrate that SPINK1 protein and mRNA levels correlate, and that low SPINK1 expression in tumor tissue is indicative of poor prognosis, in line with our previously published results (Koskensalo et al., 2012) and with a recent publication (Chen et al., 2016) in which high SPINK1 tumor expression correlated with a good prognosis in CRC patients receiving cetuximab therapy. Further, here we show for the first time that expression of BRAF V600E mutation correlates with low SPINK1 expression level. The ExBP-RT method used for the BRAF V600E analyses detects mRNA of expressed mutations in tumor tissue, rather than the presence of mutated DNA. This allowed us to use the same patient samples to analyze the expression of both SPINK1 and BRAF V600E mRNA.

Accepted Article

These clinical findings of BRAF V600E and SPINK1 expression in our CRC patient cohort led us to hypothesize that MAPK inhibitors might affect SPINK1 levels. In order to test this, we used a panel of colorectal cancer cell lines harboring the V600E mutant BRAF and compared the effects to a BRAF wild type CRC cell line. Vemurafenib treatment is not beneficial in BRAF-mutant CRC patients and combination therapies with MAPK inhibitors with EGFR inhibitors are under clinical investigations. A phase I study by Corcoran and colleagues (Corcoran et al., 2015) suggested that dual MAPK pathway blockade with the BRAF inhibitor dabrafenib and the MEK inhibitor trametinib can lead to a meaningful clinical benefit in a subset of patients with BRAF V600E metastatic CRC. Based on our results, trametinib treatment, which inhibits both MEK1 and MEK2, might be an effective therapy in BRAF V600E-positive / SPINK1-low subpopulation of CRC patients, as it led to increased SPINK1 secretion in BRAF V600E positive colorectal cancer cells. Furthermore, it was the only MAPK inhibitor that was able to diminish phosphorylation of MEK and ERK in the BRAF V600E CRC cell lines Colo205 and HT-29. Our data also supports both preclinical and clinical findings that vemurafenib is not effective in BRAF-mutant CRC, as it was not capable to completely suppress MAPK signaling. As PD98059, a MEK1 inhibitor, was not able to affect SPINK1 secretion or MEK and ERK phosphorylation, our data implicates MEK2 as a critical protein in the MAPK pathway in colorectal adenocarcinoma.

Bidirectional kinase-protease interactions are known to have a role in cancer and clinical implications of such kinase-protease crosstalk have started to emerge (Lopez-Otin and Hunter, 2010). In our study, parallel to SPINK1 increase, we observed a decrease in the expression and secretion of trypsin-1 and -2 in response to vemurafenib, trametinib and SCH772984. Studies have suggested that robust MAPK pathway suppression is required for response in BRAF V600E cancers and acquired resistance to BRAF inhibitor combinations involve reactivation of the MAPK pathway (Ahronian et al., 2015). Recently Miller and colleagues (Miller et al. 2016) showed that MEK inhibitors lead to reduced proteolytic shedding of cell-surface receptor tyrosine kinases by inhibiting the catalytic activity of A Disintegrin and Metalloproteinases (ADAM), thus leading to increased mitogenic signaling and kinase inhibitor resistance. Further, disrupting the protease inhibition by neutralizing a

putative ADAM10 inhibitor tissue inhibitor of metalloproteinases 1 (TIMP1), MAPK inhibitor efficacy was improved (Miller et al. 2016). These findings, along with ours, highlight the extensive crosstalk between kinases, proteases and cognate protease inhibitors in response to molecularly targeted therapies and warrant further studies.

Here we describe a novel mechanism of ATF-4 transcription repression by molecularly targeted therapy, as trametinib was able to downregulate ATF-4 transcription leading to reduced ATF-4 protein level in Colo205 and HT-29 cells. ATF-4 is a well-characterized effector of integrated stress response. It has several dimerization partners that influence its gene transcription, thus guiding cellular outcomes (Pakos-Zebrucka et al., 2016). In most cases cellular stress induces upregulation of ATF-4 transcription. For example in BRAF inhibitor-sensitive melanoma cell lines, the preclinical version of vemurafenib, PLX4720, led to rapid induction of ATF-4 (Ma et al., 2014). However, there is evidence of transcriptional repression of ATF-4 by some cellular stressors, such as C/EBP β during UV irradiation and in non-alcoholic fatty liver and non-alcoholic steatohepatitis (Pakos-Zebrucka et al., 2016). Of note, the mechanism by which these MAPK inhibitors activate SPINK1 transcription remains to be revealed, as on the transcription factor array we did not observe any significant increases in response to the trametinib treatment.

5. Conclusions

In conclusion, the present study demonstrates for the first time an inverse relationship between expressed BRAF V600E mutations and SPINK1 expression. Further, we show that in addition of downregulating phosphorylation of ERK and MEK, trametinib treatment leads also to downregulation of ATF-4 and trypsin 1 and -2 with concomitant increase in SPINK1 secretion. Both ATF-4 and trypsins have been shown to confer survival advantage of cancer cells and thereby to regulate tumor progression. Thus finding an effective way to inhibit the expression of these proteins while at the same time sustaining SPINK1 levels might have a clinical benefit in BRAF V600E positive colorectal adenocarcinoma. Although further studies are warranted, SPINK1 expression seems to be a useful

Molecular Oncology (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd.

biomarker in CRC and its expression might guide patient stratification and treatment response to molecularly targeted therapies.

Acknowledgements

Anne Ahmanheimo, Maarit Leinimaa and Kristiina Nokelainen are thanked for technical assistance. High Throughput Biomedicine Unit (Institute for Molecular Medicine Finland FIMM) is thanked for providing the robotics for the qPCR set-up.

Funding

Orion Research Foundation, Ruth and Nils-Erik Stenbäck foundation, Finska Läkaresällskapet, the Sigrid Jusélius Foundation and the Finnish Cancer Foundation

Author contributions

KR and JS were responsible for the study conception, design and data analysis; SL and CH were responsible for the patient identification and sample collection; KXD and THH were responsible for the ExBP-RT data acquisition; KR was responsible for the *in vitro* studies; HM was responsible for the statistical analyses. All authors were responsible for the data interpretation and manuscript writing. All authors read and approved the final version of the manuscript.

Disclosures

The authors disclose no potential conflicts of interest.

References

Ahronian L.G., Sennott E.M., Van Allen E.M., Wagle N., Kwak E.L., Faris J.E., et al., 2015. Clinical Acquired Resistance to RAF Inhibitor Combinations in BRAF-Mutant Colorectal Cancer through MAPK Pathway Alterations. *Cancer Discov.* 5, 358-367.

Barras D., 2015. *BRAF* Mutation in Colorectal Cancer: An Update. *Biomark Cancer.* 7, 9–12.

Chen Y.T., Tsao S.C., Yuan S.S., Tsai H.P., Chai C.Y., 2015. Serine Protease Inhibitor Kazal Type 1 (SPINK1) Promotes Proliferation of Colorectal Cancer Through the Epidermal Growth Factor as a Prognostic Marker. *Pathol Oncol Res.* 21, 1201-1208.

Chen Y.T., Tsao S.C., Tsai H.P., Wang J.Y., Chai C.Y., 2016. Serine protease inhibitor Kazal type 1 (SPINK1) as a prognostic marker in stage IV colon cancer patients receiving cetuximab based targeted therapy. *J Clin Pathol.* <http://dx.doi.org/10.1136/jclinpath-2016-203638>

Corcoran R.B., Atreya C.E., Falchook G.S., Kwak E.L., Ryan D.P., Bendell J.C., et al., 2015. Combined BRAF and MEK Inhibition With Dabrafenib and Trametinib in BRAF V600-Mutant Colorectal Cancer. *J Clin Oncol.* 33:4023-31.

Dienstmann R., Tabernero J., 2016. Spectrum of Gene Mutations in Colorectal Cancer: Implications for Treatment. *Cancer J.* 22, 149-155.

Elez E., Schellens J., van Geel R., Bendell J., Spreafico A., Schuler M. et al. 2015. Results of a phase 1b study of the selective BRAF V600 inhibitor encorafenib in combination with cetuximab alone or cetuximab + alpelisib for treatment of patients with advanced BRAF-mutant metastatic colorectal cancer. *AnnOncol.* 26(suppl 4):iv120. <https://doi.org/10.1093/annonc/mdv262.08>

Gaber A., Nodin B., Hotakainen K., Nilsson E., Stenman U.H., Bjartell A., et al., 2010. Increased serum levels of tumour-associated trypsin inhibitor independently predict a poor prognosis in colorectal cancer patients. *BMC Cancer.* 10:498 <http://dx.doi.org/10.1186/1471-2407-10-498>

Ho T.H., Dang K.X., Lintula S., Hotakainen K., Feng L., Olkkonen V.M., et al., 2015. Extendable blocking probe in reverse transcription for analysis of RNA variants with superior selectivity. *Nucleic Acids Res.* 43, e4. <http://dx.doi.org/10.1093/nar/gku1048>

Ida S., Ozaki N., Araki K., Hirashima K., Zaitu Y., Taki K., et al., 2015. SPINK1 Status in Colorectal Cancer, Impact on Proliferation, and Role in Colitis-Associated Cancer. *Mol Cancer Res.* 13, 1130-1138.

Itkonen O, Koivunen E, Hurme M, Alfthan H, Schröder T, Stenman UH., 1990. Time-resolved immunofluorometric assays for trypsinogen-1 and 2 in serum reveal preferential elevation of trypsinogen-2 in pancreatitis. *J Lab Clin Med.* 115:712-718.

Janeiro E., Guimarães J., Stenman U.H., Catarino M., Itkonen O., 2012. Validation and comparison of tumor-associated trypsin inhibitor (TATI) immunoassays. *Clin Chim Acta.* 413,1244-1248.

Koivunen E., Itkonen O., Halila H., Stenman U.H., 1990. Cyst fluid of ovarian cancer patients contains high concentrations of trypsinogen-2. *Cancer Res.* 50, 2375-2378.

Koskensalo S., Hagström J., Louhimo J., Stenman U.H., Haglund C., 2012. Tumour-associated trypsin inhibitor TATI is a prognostic marker in colorectal cancer. *Oncology.* 82, 234-241.

Koskensalo S., Louhimo J., Hagström J., Lundin M., Stenman U.H., Haglund C., 2013. Concomitant tumor expression of EGFR and TATI/SPINK1 associates with better prognosis in colorectal cancer. *PLoS One.* 8, e76906 <http://dx.doi.org/10.1371/journal.pone.0076906>

Livak K.J. and Schmittgen T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 25, 402-408.

Lopez-Otin C. and Hunter T., 2010. The regulatory crosstalk between kinases and proteases in cancer. *Nat Rev Cancer.* 10, 278-292.

Ma X.H., Piao S.H., Dey S., McAfee Q., Karakousis G., Villanueva J., et al., 2014. Targeting ER stress-induced autophagy overcomes BRAF inhibitor resistance in melanoma. *J Clin Invest.* 124,1406–1417.

Miller M.A., Oudin M.J., Sullivan R.J., Wang S.J., Meyer A.S., Im H. et al., 2016. Reduced Proteolytic Shedding of Receptor Tyrosine Kinases Is a Post-Translational Mechanism of Kinase

Inhibitor Resistance. *Cancer Discov.* 6, 382-399.

Moriarty A., O'Sullivan J., Kennedy J., Mehigan B., McCormick P., 2016. Current targeted therapies in the treatment of advanced colorectal cancer: a review. *Ther Adv Med Oncol.* 8, 276–293.

Osman S., Turpeinen U., Itkonen O., Stenman U.H., 1993. Optimization of a time-resolved immunofluorometric assay for tumor-associated trypsin inhibitor (TATI) using the streptavidin-biotin system. *J Immunol Methods.* 161, 97-106.

Paju A., Sorsa T., Tervahartiala T., Koivunen E., Haglund C., Leminen A., et al., 2001. The levels of trypsinogen isoenzymes in ovarian tumour cyst fluids are associated with promatrix metalloproteinase-9 but not promatrix metalloproteinase-2 activation. *Br J Cancer.* 84, 1363-1371.

Pakos-Zebrucka K., Koryga I., Mnich K., Ljujic M., Samali A., Gorman AM., 2016. The integrated stress response. *EMBO Rep.* 17, 1374-1395.

Prahalad A., Sun C., Huang S., Di Nicolantonio F., Salazar R., Zecchin D., et al., 2012. Unresponsiveness of colon cancer to BRAF(V600E) inhibition through feedback activation of EGFR. *Nature* 483, 100-103.

Räsänen K., Salmenperä P., Baumann M., Virtanen I., Vaheri A., 2008. Nemo-sis of fibroblasts is inhibited by benign HaCaT keratinocytes but promoted by malignant HaCaT cells. *Mol Oncol.* 2, 340-348.

Räsänen K., Itkonen O., Koistinen H., Stenman U.H. 2016a. Emerging Roles of SPINK1 in Cancer. *Clin Chem.* 62:449-457.

Räsänen K., Lehtinen E., Nokelainen K., Kuopio T., Hautala L., Itkonen O., et al., 2016b. Interleukin-6 increases expression of serine protease inhibitor Kazal type 1 through STAT3 in colorectal adenocarcinoma. *Mol Carcinog.* 55, 2010-2023.

Semrad T.J., Kim E.J., 2016. Molecular testing to optimize therapeutic decision making in advanced colorectal cancer. *J Gastrointest Oncol.* S11-20.

Tables

Table 1. Patient characteristics

All patients	N=571	N	%
Age	<65	231	40
	>=65	340	60
Gender	female	264	46
	male	307	54
Type	adeno	513	90
	mucinous	58	10
Location	Colon	381	67
	Rectum	190	33
Side	dex	201	35
	sin	369	65
Dukes	A	74	13
	B	211	37
	C	166	29
	D	120	21
Grade	1	28	5
	2	392	69
	3	109	19
	4	21	4
Age (min-max), years	68.1	(29.3-97.2)	

Table 2. Primer and probe sequences for qPCR step of different ExBP-RT assays (Locked Nucleic Acid (LNA) = [+A], [+G], [+C], [+T]; Inosine = i; 6-carboxyfluorescein: FAM; Black Hole Quenchers: BHQ).

Primers and probes	Sequences (5' – 3')	Concentrations
<i>Mutant BRAF V600E assays</i>		
BRAF Forward primer	5'-AGACCTCACAGTAAAAATAGGTGA-3'	0.5 μM
Common Reverse primer	5'-CGATCAGACGACGAC-3'	0.5 μM
BRAF-Probe	FAM-TTC[+T]CT[+G]TA[+G]CT[+A]GACCAA-BHQ1	0.1 μM
<i>Total-BRAF assays</i>		
Total BRAF Forward primer	5'-CATGAAGACCTCACAGTAAA-3'	1.5 μM
Total BRAF Reverse primer	5'-GATTTCACTGTAGCTAGACC-3'	1.5 μM

Table 3. Statistical analyses of expressed BRAF V600E mutations and SPINK1 expression

		HR	95% CI		p
<i>Univariate</i>			lower	upper	
Age 65 yr	Age=> 65 vs <65	1.45	1.11	1.90	0.006
Gender	male vs female	1.03	0.80	1.34	0.807
Dukes	A	1.00			
	B	2.09	0.98	4.45	0.056
	C	6.52	3.15	13.49	0.000
	D	27.92	13.48	57.84	0.000
Grade	1	1.00			
	2	1.96	0.87	4.42	0.106
	3	2.88	1.24	6.71	0.014
	4	3.09	1.12	8.50	0.029
Location	rectum vs colon	1.25	0.96	1.64	0.095
Side	sin vs dex	1.25	0.95	1.66	0.116
Type	mucinous vs adeno	0.93	0.61	1.42	0.744
BRAF V600E mutation	yes vs no	2.12	1.48	3.03	0.000
SPINK1 mRNA	Continuous	0.93	0.85	1.01	0.098
SPINK1 mRNA dichotomous	>2.4 vs <=2.4	0.556	0.321	0.961	0.036
SPINK1 mRNA	<0.5	1			
	0.5-2.4	0.87	0.63	1.20	0.403
	>=2.4	0.51	0.29	0.91	0.023
BRAF V600E and SPINK1 mRNA	BRAF V600E mutation	1.00			
	SPINK1<=2.4, no mutation in BRAF	0.40	0.26	0.62	0.000
	SPINK1>2.4, no mutation in BRAF	0.23	0.12	0.45	0.000
<i>Multivariate BRAF</i>					
Age 65 yr	Age=> 65 vs <65	2.01	1.52	2.66	0.000
Gender	male vs female	1.14	0.87	1.48	0.344
Dukes	A	1			
	B	2.17	1.01	4.66	0.047
	C	7.08	3.39	14.76	0.000
	D	32.16	15.36	67.33	0.000
Location	rectum vs colon	1.47	1.12	1.93	0.006
BRAF V600E mutation	yes vs no	2.84	1.84	4.40	0.000
BRAF V600E time dependent	After two years of survival	0.16	0.05	0.54	0.003
<i>Multivariate SPINK1</i>					
Age 65 yr	Age=> 65 vs <65	2.20	1.58	3.07	0.000
Gender	male vs female	1.12	0.81	1.54	0.489
Dukes	A	1			
Dukes_B	B	2.68	1.02	7.02	0.045
Dukes_C	C	8.71	3.44	22.06	0.000
Dukes_D	D	40.93	16.03	104.52	0.000
Location	rectum vs colon	1.66	1.19	2.32	0.003
SPINK1 mRNA	<0.5	1			
	0.5-2.4	0.72	0.50	1.02	0.061
	>=2.4	0.43	0.22	0.84	0.014
SPINK1 mRNA time dependent	After five years of survival	2.48	1.13	5.41	0.023
<i>Multivariate SPINK1 and BRAF</i>					
Age 65 yr	Age=> 65 vs <65	2.12	1.51	2.96	0.000
Gender	male vs female	1.13	0.82	1.55	0.465
Dukes	B	2.77	1.06	7.28	0.038

	C	8.79	3.47	22.26	0.000
	D	43.59	17.02	111.65	0.000
Location	rectum vs colon	1.75	1.25	2.45	0.001
BRAF V600E and SPINK1 mRNA	BRAF V600E mutation	1			
	SPINK1 \leq 2.4, no mutation in BRAF	0.29	0.17	0.47	0.000
	SPINK1 $>$ 2.4, no mutation in BRAF	0.09	0.03	0.27	0.000
BRAF V600E and SPINK1 mRNA	After two years of survival	3.49	1.43	8.54	0.006

Table 4. Basal levels of secreted SPINK1, trypsin-1 and -2 at 72h time point.

<i>Cancer type</i>	<i>Cell line</i>	<i>BRAF status</i>	<i>SPINK1 ng/ml</i>	<i>Trypsin-1 ng/ml</i>	<i>Trypsin-2 ng/ml</i>
CRC	Colo205	V600E	1.6 (\pm 0.06)	12.2 (\pm 0.8)	56 (\pm 6.2)
CRC	HT-29	V600E	11.4 (\pm 2.4)	2.5 (\pm 0.3)	3.9 (\pm 0.5)
CRC	Caco-2	WT	35.3 (\pm 4.3)	1.3 (\pm 0.3)	ND
CRC	SW480	WT	ND	ND	ND
Melanoma	SK-MEL-2	WT	ND	ND	ND
Melanoma	SK-MEL-5	V600E	ND	ND	ND

Figure captions

Figure 1. Survival curves for BRAF wild-type and V600E expressing colorectal cancer patients.

Figure 2. (A) Representative IHC images of SPINK1 a) negative, b) positive ($<$ 2.4) and c) positive ($>$ 2.4) immunoexpression. Images taken at 10x magnification. The value below is the relative SPINK1 mRNA expression in the same sample calculated using the $\Delta\Delta C_t$ method. (B) Survival curves for SPINK1 mRNA ($<$ 2.4) and ($>$ 2.4) colorectal cancer patients. (C) Survival curves for SPINK1 mRNA ($<$ 2.4) and ($>$ 2.4) colorectal cancer patients having disease on the left side.

Figure 3. MAPK pathway inhibitors increase SPINK1 levels in BRAF V600E colorectal cancer. Secreted SPINK1 protein levels were analyzed by IFMA in Colo205 (A), HT-29 (B) and (C) Caco-2 cells at 72h timepoint. Vemurafenib, trametinib and SCH772984 increased SPINK1 secretion in Colo205 and HT-29 cells over 2-fold (dashed line), whereas no 2-fold increase was seen in the Caco-2 cells compared to control and DMSO-treated cells.

Figure 4. Time-dependent increase of SPINK1 levels in response to vemurafenib, trametinib and SCH772984 in Colo205 and HT-29 cells. (A) In Colo205 cells vemurafenib, trametinib and SCH772984 significantly increased SPINK1 secretion at 48 and 72 hour time points with concomitant decrease in trypsin-1 and -2 secretion at 72h timepoint as measured by respective IFMAs. (B) In HT-29 cells vemurafenib, trametinib and SCH772984 significantly increased SPINK1 secretion at 48 and 72 hour time points. Vemurafenib and SCH772984, but no trametinib, reduced trypsin-1 and -2 levels in HT-29 cells at 72h timepoint. (C) Vemurafenib, trametinib and SCH772984 did not increase SPINK1 secretion or decrease trypsin-1 secretion in Caco-2 cells compared to control or DMSO-treated cells. Significantly different (* $p < 0.05$) as compared to control by two tailed t-test. Trypsin-2 was not detected in HT-29 cells at 24h timepoint and in Caco-2 cells at any timepoint.

Figure 5. Trametinib diminishes ERK and MEK phosphorylation in BRAF V600E CRC cells. ERK1/2 (Thr202/Tyr204) and MEK1/2 (Ser217/221) residues are dephosphorylated by trametinib (60 nM) as shown by a Western blot of whole cell lysates of Colo205 and HT-29 cell lines at 24h timepoint. Vemurafenib and SCH772984 reduce ERK1/2 (Thr202/Tyr204) and MEK1/2 (Ser217/221) phosphorylation to a lesser extent in Colo205 and HT-29 cells. Vemurafenib (Vem.), trametinib (Tram.), SCH772984 (SCH) or PD98059 (PD). Total ERK1/2 and MEK1/2 antibodies were used as controls.

Figure 6. Trametinib downregulates ATF-4 mRNA and protein levels. (A) Colo205 and HT-29 cells were treated with 60 nM trametinib for 24h after which RNA was extracted and TaqMan non-HOX transcription factor array was performed. ATF-4 mRNA was decreased by two-fold in response to trametinib when compared to DMSO control. (B) Western blot of whole cell lysates of Colo205 and HT-29 cells harvested after 24h treatment with either 60 nM vemurafenib (Vem.), trametinib (Tram.), SCH772984 (SCH) or PD98059 (PD) showing downregulation of ATF-4 protein level in response to vemurafenib. GAPDH was used as a loading control.

Supplementary Figure 1. (A) *SPINK1*, (B) *PRSS1* and (C) *PRSS2* mRNA levels analyzed by qPCR in response to inhibitor treatment (60 nM) at 72h time point. Vemurafenib, trametinib and SCH772984 increase *SPINK1* mRNA expression in Colo205 and HT-29 cells. In Colo205 cells vemurafenib, trametinib and SCH772984 lead to concomitant decrease in *PRSS1* and *PRSS2* mRNA expression; PD98059 also downregulates the expression of *PRSS1* and *PRSS2* in Colo205 cells. In HT-29 cells vemurafenib decreases *PRSS1* and *PRSS2* expression, SCH772984 decreases *PRSS1* expression and trametinib increases the expression of *PRSS1* and *PRSS2*. In Caco-2 cells *SPINK1* expression is slightly increased in response to trametinib and SCH77298; *PRSS1* or *PRSS2* mRNA levels are not affected by MAPK inhibitors. Vem.: vemurafenib; Tram.: trametinib; SCH: SCH772984; PD: PD98059

Supplementary Figure 2. Western blot of whole cell lysates of Colo205 and HT-29 cells harvested after 24h treatment with either 60 nM vemurafenib (Vem.), trametinib (Tram.), SCH772984 (SCH) or PD98059 (PD). STAT3 protein levels remained constant in response to MAPK inhibitors. GAPDH was used as a loading control









