


Journal Pre-proof



Identification of required host factors for SARS-CoV-2 infection in human cells

Zharko Daniloski, Tristan X. Jordan, Hans-Hermann Wessels, Daisy A. Hoagland, Silva Kasela, Mateusz Legut, Silas Maniatis, Eleni P. Mimitou, Lu Lu, Evan Geller, Oded Danziger, Brad R. Rosenberg, Hemali Phatnani, Peter Smibert, Tuuli Lappalainen, Benjamin R. tenOever, Neville E. Sanjana

PII: S0092-8674(20)31394-5

DOI: <https://doi.org/10.1016/j.cell.2020.10.030>

Reference: CELL 11695

To appear in: *Cell*

Received Date: 12 August 2020

Revised Date: 25 September 2020

Accepted Date: 20 October 2020

Please cite this article as: Daniloski, Z., Jordan, T.X., Wessels, H.-H., Hoagland, D.A., Kasela, S., Legut, M., Maniatis, S., Mimitou, E.P., Lu, L., Geller, E., Danziger, O., Rosenberg, B.R., Phatnani, H., Smibert, P., Lappalainen, T., tenOever, B.R., Sanjana, N.E., Identification of required host factors for SARS-CoV-2 infection in human cells, *Cell* (2020), doi: <https://doi.org/10.1016/j.cell.2020.10.030>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Inc.

*Article***Identification of required host factors for SARS-CoV-2 infection
in human cells**

Zharko Daniloski^{1,2,7}, Tristan X. Jordan^{3,7}, Hans-Hermann Wessels^{1,2}, Daisy A. Hoagland^{3,7},
Silva Kasela^{1,4}, Mateusz Legut^{1,2}, Silas Maniatis¹, Eleni P. Mimitou⁵, Lu Lu^{1,2}, Evan Geller^{1,2},
Oded Danziger³, Brad R. Rosenberg³, Hemali Phatnani^{1,6}, Peter Smibert⁵, Tuuli Lappalainen^{1,4},
Benjamin R. tenOever^{3,*}, Neville E. Sanjana^{1,2,8,*}

1. New York Genome Center, New York, NY, USA
 2. Department of Biology, New York University, New York, NY, USA
 3. Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA
 4. Department of Systems Biology, Columbia University, New York, NY, USA
 5. Technology Innovation Lab, New York Genome Center, New York, NY, USA
 6. Department of Neurology, Columbia University Irving Medical Center, New York, NY, USA
 7. These authors contributed equally
 8. Lead Contact
- * Correspondence: benjamin.tenoever@mssm.edu (B.R.t.), neville@sanjanalab.org (N.E.S.)

Keywords: COVID-19, SARS-CoV-2, genome-wide screen, CRISPR, loss-of-function, human lung

Summary

To better understand host-virus genetic dependencies and find potential therapeutic targets for COVID-19, we performed a genome-scale CRISPR loss-of-function screen to identify host factors required for SARS-CoV-2 viral infection of human alveolar epithelial cells. Top-ranked genes cluster into distinct pathways, including the vacuolar ATPase proton pump, Retromer, and Commander complexes. We validate these gene targets using several orthogonal methods such as CRISPR knock-out, RNA interference knock-down, and small-molecule inhibitors. Using single-cell RNA-sequencing, we identify shared transcriptional changes in cholesterol biosynthesis upon loss of top-ranked genes. In addition, given the key role of the ACE2 receptor in the early stages of viral entry, we show that loss of *RAB7A* reduces viral entry by sequestering the ACE2 receptor inside cells. Overall, this work provides a genome-scale, quantitative resource of the impact of the loss of each host gene on fitness/response to viral infection.

* * *

1 **Introduction**

2 As of October 2020, SARS-CoV-2 (Severe Acute Respiratory Syndrome-Coronavirus-2), the
3 virus that causes COVID-19, has infected 40 million people worldwide and led to the deaths of
4 more than 1 million people, according to the John Hopkins Research Center (Gardner, 2020).
5 SARS-CoV-2 belongs to the family of enveloped viruses known as *Coronaviridae* and was first
6 reported in late 2019 in China. Over the past two decades, it is the third zoonotic coronavirus to
7 emerge: Compared to the other two coronaviruses, SARS-CoV (2002) and Middle East
8 Respiratory Syndrome (MERS)-CoV (2012), SARS-CoV-2 shows an increased infectivity and
9 lower case-fatality rate, contributing to its wide-spread transmission and resulting in a pandemic
10 (Gates, 2020; Liu et al., 2020). Given that SARS-CoV-2 has already taken a major toll on human
11 life and livelihoods worldwide, many research institutions, governmental organizations and
12 pharmaceutical companies are working to identify antiviral drugs and develop vaccines.
13 Currently, there are nearly 30 vaccines against SARS-CoV-2 in clinical trials and an FDA
14 approved antiviral drug (remdesivir) that acts as an inhibitor of the SARS-CoV-2 viral RNA-
15 dependent RNA polymerase (Beigel et al., 2020; Funk et al., 2020). A recent study identified
16 small molecules that antagonize SARS-CoV-2 replication and infection by testing ~12,000
17 clinical-stage and FDA-approved inhibitors (Riva et al., 2020). Here, we utilize an alternative
18 approach — a genome-scale loss-of-function screen — to identify targets among host genes that
19 are required for SARS-CoV-2 infection. These gene targets (and inhibitors of these genes) may
20 aid in the development of new therapies for COVID-19.

21
22 SARS-CoV-2 is an enveloped positive-sense RNA virus that relies on host factors for all stages
23 of its lifecycle (Kim et al., 2020; Zhou et al., 2020). The viral envelope is coated by Spike
24 protein trimers that bind to angiotensin converting-enzyme 2 (ACE2) receptor, which is required
25 for SARS-CoV-2 infection (Hoffmann et al., 2020a; Zhou et al., 2020). The Spike protein
26 undergoes proteolytic cleavage that is catalyzed by several host proteases, such as furin,
27 TMPRSS2 and Cathepsin L, and can occur in the secretory pathway of the host cell or during
28 viral entry in the target cell. Proteolytic cleavage is considered to be required for activation of
29 Spike that in turn allows for viral-host membrane fusion and release of the viral RNA into the
30 host cytoplasm (Hoffmann et al., 2020b). Once in the cytoplasm, the virus utilizes the host and
31 its own machinery to replicate its genetic material and assemble new viral particles. Recent

32 proteomic studies have identified hundreds of host proteins that directly bind to SARS-CoV-2
33 viral proteins and have mapped changes in the global protein phosphorylation landscape in
34 response to viral infection, highlighting the interest in better understanding of host-virus genetic
35 dependencies (Bouhaddou et al., 2020; Gordon et al., 2020). To date, there are no genome-wide
36 studies that directly identify human genes required for viral infection, which will be of great
37 interest and utility for the broader scientific community.

38
39 Here, we perform a genome-scale CRISPR loss-of-function screen in human alveolar basal
40 epithelial carcinoma cells to identify genes whose loss confers resistance to SARS-CoV-2 viral
41 infection. We validate that these genes reduce SARS-CoV-2 infection using multiple orthogonal
42 cell perturbations (CRISPR knock-out, RNA interference knock-down, and small-molecule
43 inhibitors). For the top gene hits, we explore potential mechanisms of their antiviral activity
44 using single-cell transcriptomics, flow cytometry and immunofluorescence. Using single-cell
45 transcriptomics we identified a group of genes (*ATP6AP1*, *ATP6VIA*, *NPC1*, *RAB7A*, *CCDC22*,
46 and *PIK3C3*) whose knockout induced shared transcriptional changes in cholesterol biosynthesis
47 pathway. Perturbation of the cholesterol biosynthesis pathway with the small molecule
48 amlodipine reduced viral infection. In addition, we show that loss of *RAB7A* reduces viral entry
49 by sequestering ACE2 receptors inside cells through altered endosomal trafficking. Prior to this
50 study, our knowledge of essential host genes for SARS-CoV-2 has been limited to only a handful
51 of genes, such as ACE2 and cathepsin L: This work provides a quantitative resource of the
52 impact of each gene's loss on response to viral infection for every protein-coding gene in the
53 human genome.

54

55 **Results**

56 *A high-throughput screen to identify genes required for SARS-CoV-2 infection*

57 To identify key genes required for SARS-CoV-2 infection, we performed a genome-scale loss-
58 of-function screen targeting 19,050 genes in the human genome using the GeCKOv2 CRISPR-
59 Cas9 library (Sanjana et al., 2014). The GeCKOv2 library contains 122,411 CRISPR single-
60 guide RNAs (6 guide RNAs per gene) and has previously been used in CRISPR screens for drug
61 resistance, immunotherapy, synthetic lethality, mitochondrial disease, and therapeutic discovery
62 for muscular dystrophy (Erb et al., 2017; Jain et al., 2016; Lek et al., 2020; Patel et al., 2017;

63 Shalem et al., 2014). First, we transduced a human alveolar basal epithelial carcinoma cell line
64 (A549) that constitutively expresses ACE2 (referred to as A549^{ACE2}) with an all-in-one lentiviral
65 vector containing Cas9, guide RNAs from the GeCKOv2 human library, and a puromycin
66 resistance gene. The transduction was performed at a low multiplicity of infection (MOI ~ 0.2) to
67 ensure that most cells would receive only one guide RNA construct (**Figure 1A**). We then
68 selected with puromycin so that only library-transduced cells remained. We also measured the
69 survival rate after puromycin selection was complete (3 days) to ensure high coverage of the
70 122,411 guide RNAs (~1,000 cells per guide RNA). After puromycin selection was complete,
71 we cultured the cells for 9 days to ensure protein depletion after CRISPR gene targeting.

72
73 Next, we infected the GeCKOv2 pool of A549^{ACE2} cells with SARS-CoV-2 virus (Isolate USA-
74 WA1/ 2020 NR-52281) at either a high (0.3) or a low (0.01) MOI. We verified that SARS-CoV-
75 2 infects A549^{ACE2} cells by staining for the nucleocapsid (N) protein at 24 hours post-infection
76 (**Figure 1B**) and, at day 6 post-infection, we measured cell survival for both the high and low
77 MOI conditions (**Figure 1C**). As expected, the higher MOI infection resulted in fewer surviving
78 cells at day 6 post-infection. Next, we extracted genomic DNA and, via amplicon sequencing, we
79 quantified guide abundance in each biological condition (**Figure 1A**). To confirm that library
80 representation was properly maintained, we computed the correlation between the guide
81 representation in the plasmid library and after puromycin selection ($r = 0.84$) (**Figure S1A**). In
82 contrast, after SARS-CoV-2 infection, there was a much greater degree of guide dropout, as
83 expected given that SARS-CoV-2 rapidly kills A549^{ACE2} cells without CRISPR perturbations
84 (**Figures 1C, D**).

85
86 Using robust-rank aggregation (RRA) on the guide relative enrichments, we computed gene-
87 level scores to identify genes where loss-of-function mutations led to enrichment within the pool
88 (**Figure 1E**) (Kolde et al., 2012). We identified approximately 1,000 genes with significant RRA
89 enrichment ($p < 0.05$) (**Figure S1B**). We also used two other previously published methods to
90 compute gene enrichments (RIGER weighted-sum and second-best rank) and found a high
91 degree of overlap between enriched genes identified by all three methods (**Figure S1C**) (Chen et
92 al., 2015; Luo et al., 2008). We also found a high degree of shared genes across both the low and
93 high SARS-CoV-2 MOI conditions: When examining the top 50 most enriched genes, we found

94 that 27 of them were shared between the low and high MOI conditions (**Figure 1F, Table S1**),
95 suggesting that several host genes involved in viral pathogenesis function independently of viral
96 dose. An independent genome-scale CRISPR screen for SARS-CoV-2 infection also performed
97 in A549 that overexpress ACE2 but with a different CRISPR library identified similar top-
98 ranked genes (Zhu et al., 2020), highlighting the robustness of our results (**Figure S1D**).

99 *Enriched genes are involved in multiple aspects of the viral lifecycle and are broadly expressed*

100 Upon close examination of the most enriched genes, we found genes involved in key aspects of
101 viral entry and replication (**Figure 2**) (Du et al., 2009). For example, the well-established entry
102 receptor angiotensin-converting enzyme 2 (ACE2) receptor was ranked as the 8th most-enriched
103 gene in the low MOI screen and 12th in the high MOI screen (**Table S1**) (Hoffmann et al., 2020a;
104 Zhou et al., 2020). Among the top 50 enriched genes, we identified several sets of related genes
105 that function together in complexes, giving us further confidence in the genome-scale screen
106 (**Figures 2 and 3A**). We found genes essential for initial attachment and endocytosis (*ACE2*,
107 *RAB7A* and 4 members of the ARP2/3 complex: *ACTR2*, *ACTR3*, *ARPC3*, and *ARPC4*), Spike
108 protein cleavage and viral membrane fusion (*CTSL* and 13 members of the vacuolar-ATPase
109 proton pump: *ATP6AP1*, *ATP6AP2*, *ATP6V0B*, *ATP6V0C*, *ATP6V0D1*, *ATP6VIA*, *ATP6VIB2*,
110 *ATP6VIC1*, *ATP6VIE1*, *ATP6VIG1*, *ATP6VIH*, *TMEM199*, and *TORIAIP1*), endosome
111 recycling (4 members of the endosomal protein sorting Retromer complex: *VPS26A*, *VPS29*,
112 *VPS35*, and *SNX27*; 4 members of the endosomal trafficking Commander complex: *COMMD2*,
113 *COMMD3*, *COMMD3-BMI1*, and *COMMD4*; and 3 members of the PI3K pathway:
114 *PIK3C3/VPS34*, *WDR81*, and *ACP5*), ER-Golgi trafficking (*DPM3*, *ERMP1*, *PPID*, and
115 *CHST14*), and transcriptional modulators (*SLTM* and *SPEN*). A consistent theme among the
116 enriched complexes is endosome function and regulation (V-ATPase proton pump, Retromer,
117 Commander, Class 3 PI3Ks) (Banerjee and Kane, 2020; Mallam and Marcotte, 2017; McNally
118 and Cullen, 2018). Gene Set Enrichment Analysis on the full ranked list of genes identified
119 significantly-enriched Gene Ontology (GO) categories for endosome processing, transport and
120 acidification and categories related to cytokinesis and virion attachment (FDR $q < 0.1$) (**Figures**
121 **3B and S2A-D**) (**Table S2**) (Subramanian et al., 2005).

122

123 Although we performed our CRISPR screen in human lung cells, we explored whether the
124 expression of host genes whose loss reduces SARS-CoV-2 infection were lung-specific or more
125 broadly expressed. To answer this question, we took the top-ranked genes and examined their
126 expression across 12 tissues using 4,790 RNA-sequencing datasets from the Genotype-Tissue
127 Expression (GTEx) v8 database (**Figure 3C**) (Aguet et al., 2019). Virtually all of the top gene
128 hits were broadly expressed across all tissues, implying that these mechanisms may function
129 independent of cell or tissue type. Among the top-ranked genes, only ACE2 showed tissue-
130 specific expression with a particular enrichment in testis, small intestine, kidney and heart
131 (**Figure 3C**).

132 *Enriched genes have been suggested to interact with viral proteins and are also essential for*
133 *other viral pathogens*

134 Recently, Krogan and colleagues performed an in-depth study of SARS-CoV-2 protein-protein
135 interaction networks by over-expressing affinity-tagged versions of each protein encoded in the
136 viral genome followed by tandem mass spectrometry after pulldown (Gordon et al., 2020). Their
137 study identified 332 high-confidence SARS-CoV-2-human protein-protein interactions (PPIs).
138 We found that some of the highly-ranked genes from our loss-of-function screen were reported
139 to have direct PPI with different viral proteins (**Figure 3D, Table S2**). For example, two highly-
140 ranked subunits of the vacuolar-ATPase proton pump, *ATP6API* and *ATP6VIA*, interact with
141 SARS-CoV-2 non-structural protein 6 (nsp6) and membrane (M) protein, respectively.
142 *ATP6API*, which was ranked 2nd in the low MOI CRISPR screen and 4th in the high MOI
143 CRISPR screen, has a very strong PPI interaction with nsp6 (Mass spectrometry Interaction
144 Statistics [MIST] score = 0.99) (Verschueren et al., 2015). Another key endocytosis protein,
145 *RAB7A*, is ranked in the top 50 genes in both CRISPR screens and interacts strongly with non-
146 structural protein 7 (nsp7) (MIST score = 0.97). We also compared the top-ranked genes with
147 another proteomic study that used proximity labeling in A549 cells over-expressing BioID-
148 tagged viral proteins and found that 22 out of the top 50 low MOI CRISPR screen genes had
149 direct interactions with viral genes — a significant enrichment over randomly chosen genes ($p =$
150 2×10^{-4}) (Samavarchi-Tehrani et al., 2020).

151
152 Since similar loss-of-function CRISPR screens have been performed to identify host genes
153 required for other viral pathogens, we next sought to understand whether the hits identified in our

154 SARS-CoV-2 screen were shared with those identified in prior screens for Zika virus (ZIKV)
155 and pandemic H1N1 avian influenza (IAV) (Y. Li et al. 2019, B. Li et al., 2020). We examined
156 whether top-ranked hits from the ZIKV and IAV screens shared similar genes and similar
157 functional categories. Overall, there was greater similarity between GO categories of enriched
158 genes for SARS-CoV-2 and ZIKV (**Figure 3E**) (**Table S2**). When examining the top 50 genes
159 from the SARS-CoV-2 screen, we found several genes that were highly enriched in all 3 viral
160 pathogen screens (**Figure S2E**). This group included subunits of the vacuolar-ATPase proton
161 pump, a well-known family of genes essential for acidification and endosomal processing
162 (Banerjee and Kane, 2020).

163 *Validation of enriched genes using CRISPR knock-out, RNA interference and small molecule* 164 *inhibitors*

165 To test the ability of top-ranked genes to block SARS-CoV-2 viral infection, we picked 30
166 genes ranked among the top 200 genes in our RRA analysis for independent validation. Each
167 gene was targeted with 3 guide RNAs distinct from the guides present in the GeCKOv2 genome-
168 wide CRISPR library (**Table S3**). The guides were synthesized individually and cloned into the
169 pCC1 lentiviral vector, a modified version of lentiCRISPRv2 with the F+E optimized guide
170 RNA scaffold (Legut et al., 2020). For a subset of genes, we validated Cas9-mediated loss-of-
171 function by western blot for protein expression (**Figure S3A**).

172
173 Cas9-perturbed A549^{ACE2} lines were infected with SARS-CoV-2 at an MOI of 0.1 and the
174 percentage of infected cells was determined by immunofluorescence against the SARS-CoV-2 N
175 protein at 36 hours post-infection (hpi). For all of the Cas9-perturbed cell lines we observed a
176 reduced percentage of infected cells with an up to 10-fold reduction in SARS-CoV-2 infection,
177 compared to the cell lines with non-targeting sgRNAs (**Figures 4A, B**). Among the genes where
178 loss provided the greatest protection against SARS-CoV-2 infection were vesicular trafficking
179 genes like *RAB7A*, *CCDC22* and *VPS35*, and other genes with well-established roles such as the
180 ACE2 receptor and the protease cathepsin L (CTSL) (**Figure 4B**). We found a significant
181 negative correlation between the percent infection in the arrayed CRISPR validation and the
182 median fold-change from the genome-wide CRISPR screen ($r_s = -0.6$, $p = 5 \times 10^{-4}$) (**Figure 4C**).

183

184 For these genes, we performed a full multi-step viral replication growth curve and found that the
185 CRISPR perturbations decrease viral load across all timepoints (5, 10, 24 and 48 hours)
186 compared to the non-targeting control (**Figure S3B**). To test if the identified hits block viral
187 infection in other cell lines, for 8 genes we generated polyclonal CRISPR knockouts in a human
188 liver cell line (Huh7.5^{ACE2}), which were then infected with SARS-CoV-2 at an MOI of 0.1.
189 Compared to non-targeting guide RNA controls, we found reduced infection with all 8 genes
190 tested in the Huh7.5^{ACE2} cells (**Figure S3C**).

191
192 As an orthogonal gene perturbation method, we also validated a more extensive list of top-
193 ranked genes via siRNA knockdown for 48 hours, followed by infection with SARS-CoV-2 at
194 MOI of 0.1 (**Table S3**). Quantification of N-protein immunofluorescent images revealed a
195 substantial reduction of the percent infected cells (**Figure S3D**).

196
197 By cross-referencing highly-ranked genes from the CRISPR screen with the Drug Gene
198 Interaction database (DGIdb), we identified a set of 69 druggable genes (**Figure 4D**) (**Table S4**)
199 (Cotto et al., 2018). We selected 9 genes that were a primary or a secondary target of one of the
200 26 small molecule inhibitors (**Figure 4E**). Among the 26 inhibitors, 9 are FDA approved and 7
201 are in Phase 2 or Phase 3 clinical trials for diverse diseases (**Table S4**). We pre-treated A549^{ACE2}
202 cells for 2 hours with 10 μ M of each inhibitor and then infected with SARS-CoV-2 and analyzed
203 the cells at 36 hpi. As a positive control, we also included remdesivir, which inhibits the viral
204 RNA polymerase and is the only currently approved treatment for COVID-19 in the United
205 States (Beigel et al., 2020). We assessed the efficacy of each inhibitor on blocking viral infection
206 using immunofluorescence and quantitative PCR (qPCR).

207
208 Seven of the 26 inhibitors that we tested (PIK-III, Compound-19, SAR405, autophinib, ALLN,
209 tamoxifen and ilomastat) resulted in >100-fold reduction of viral load as measured by qPCR
210 (**Figure 4E**). Among the best performing inhibitors, 4 of them target the same gene *PIK3C3*
211 (also known as *VPS34*) and 2 inhibitors (autophinib and ALLN) reduced the viral load more than
212 1000-fold. Similar results were obtained by immunofluorescent imaging of SARS-CoV-2 N
213 protein (**Figure S4A**). By testing the top four *PIK3C3* inhibitors in combination with CRISPR
214 targeting of *PIK3C3*, we found that Compound-19, PIK-III and autophinib were specific while

215 SAR405 resulted in greater viral inhibition in *PIK3C3* CRISPR-perturbed cells, suggesting some
216 potential off-target activity for SAR405 (**Figure S4B**).

217
218 Combinations of some of the top performing inhibitors overall showed an additive effect and
219 further protected the cells from SARS-CoV-2 infection (**Figure S4C**). Next we measured cell
220 viability at 36 hours post inhibitor treatment using flow cytometry. We observed more than 50%
221 reduction of A549^{ACE2} cell viability with two pan-HDAC inhibitors, panobinostat and pracinostat
222 (**Figure 4F**).

223 *Single-cell sequencing identifies cholesterol biosynthesis as a common mechanism underlying*
224 *multiple enriched genes from the CRISPR screen*

225 Next, we sought to understand the mechanisms underlying how individual genes identified in our
226 loss-of-function screen prevent SARS-CoV-2 infection and if host gene loss alters cell
227 transcriptional programs. For this, we utilized the Expanded CRISPR-compatible Cellular
228 Indexing of Transcriptomes and Epitopes by sequencing (ECCITE-seq) method to couple pooled
229 CRISPR perturbations of our top hit genes with a single-cell transcriptomic and proteomic
230 readout (Mimitou et al., 2019) (**Figure 5A**) (**Table S5**). ECCITE-seq is a high-throughput
231 approach to identify the molecular mechanisms and cellular pathways that drive infection
232 resistance; the pooled format also provides a more controlled experiment that may be less
233 susceptible to batch variation. Importantly, cells were infected at low MOI to maximize the
234 fraction of cells that express a single guide RNA, and therefore can be assigned a specific gene
235 perturbation. For this, we pooled all individual guide RNA plasmids (3 per target gene) used to
236 validate our genome-scale screen and 9 non-targeting (NT) sgRNAs (**Figure 4B**) (**Table S3**).

237
238 In an initial ECCITE-seq experiment, we identified a median of 152 single cells per target gene.
239 We observed specific reduction of target gene expression in cells grouped by target gene,
240 indicating nonsense-mediated decay of transcripts with frameshift indel mutations after CRISPR
241 modification (**Figure S5A**). This effect was more pronounced for genes with higher expression.
242 Using differential gene expression analysis between cells with non-targeting guide RNAs and
243 cells with targeting guide RNAs, we identified for 11 of the 30 target genes more than 5
244 differentially expressed genes with a minimal log fold change (see *Methods*), implying that loss
245 of these genes results in a detectable transcriptomic shift. It is likely that loss of the other 19

246 genes results in more subtle changes in only a few genes; however, even for ‘non-perturbed’
247 genes with sufficient basal expression, we detected clear changes in the CRISPR target gene. We
248 repeated the ECCITE-seq experiment focusing on the 11 genes with detectable transcriptomic
249 shifts upon target gene perturbation. Combining both replicate experiments, we obtained 18,853
250 cells that expressed only one guide RNA with a median of 1,388 cells per target gene.

251
252 We found that loss of 6 of the ‘perturbed’ genes — *ATP6AP1*, *ATP6V1A*, *CCDC22*, *NPC1*,
253 *PIK3C3* and *RAB7A*, which are part of the endosomal entry pathway — yielded similar gene
254 expression signatures among upregulated differentially expressed genes (**Figures 5B, C** and
255 **S5B**). These 6 target gene perturbations all led to upregulation of pathways affecting lipid and
256 cholesterol homeostasis (**Figure 5C**). Recently, we performed a large survey of >20,000
257 potential pharmacological treatments for COVID-19 and, for compounds effective at preventing
258 viral infection, we identified induction of the cholesterol biosynthesis pathway as a potential
259 mechanism of viral inhibition (Hoagland et al., 2020). Loss-of-function mutations in these 6
260 genes may function through a similar mechanism (induction of cholesterol synthesis) that
261 combats the virus-mediated suppression of cholesterol synthesis. Among the significant
262 differentially expressed genes, we also found 61 genes from the enriched CRISPR screen genes
263 ($n = 20$ genes up-regulated; $n = 41$ genes down-regulated) (**Table S5**). For example, *NPC1*,
264 *ATP6V1F* and *ATP6V1G1* are upregulated in most of the 6 endosomal entry pathway gene-
265 perturbed cells (**Figure 5C**), suggesting compensatory upregulation of related genes to mitigate
266 target gene loss.

267
268 To understand how these changes impact lipid production, we measured cholesterol levels in
269 cells after CRISPR perturbation and found that loss of these genes increases cholesterol by
270 between 10 and 50%, depending on the perturbation (**Figure 5D**). To show that increases in
271 cholesterol leads to increased SARS-CoV-2 resistance, we treated A549^{ACE2} cells with
272 amlodipine, a calcium-channel antagonist that increases intracellular cholesterol (Mori et al.,
273 1988; Ranganathan et al., 1982). We verified that amlodipine increases cholesterol levels in
274 A549^{ACE2} cells (**Figure S5C**) and found that pre-treatment with amlodipine results in reduced
275 SARS-CoV-2 viral infection, as measured by qPCR for nucleocapsid RNA, plaque formation
276 and number of viral RNA reads from RNA-sequencing, with only a modest impact on cell

277 viability (**Figures S5D-G**). RNA-sequencing of cells treated with amlodipine shows a similar
278 differential gene expression profile as seen in our ECCITE-seq with CRISPR perturbations with
279 the most significant upregulated pathway as cholesterol biosynthesis (**Figures S5H, I**).

280

281

Journal Pre-proof

282 *RAB7A* knockout results in a reduced cell surface expression and an increased endosomal
283 accumulation of ACE2

284 Next, we sought to determine if any of the top-ranked genes regulate cell surface expression of
285 ACE2, as surface ACE2 is required for SARS-CoV-2 infection (Hoffmann et al., 2020a). To
286 measure the cell surface expression of ACE2, we performed flow cytometry on A549 wild-type
287 cells and a panel of A549^{ACE2} CRISPR-perturbed cells. ACE2 expression was detected in
288 A549^{ACE2}, but not in A549 wild type cells, validating the antibody specificity (**Figure 6A** and
289 **S6**). Flow cytometry analysis of the A549^{ACE2} CRISPR-perturbed lines revealed that *RAB7A*
290 knock-out cells have a significantly reduced cell surface expression of ACE2 compared to cells
291 transduced with a non-targeting guide RNA (**Figure 6B**). Efficient Rab7a protein depletion
292 across the *RAB7A* CRISPR-perturbed lines used was confirmed by western blot (**Figure S6C**).

293
294 Rab7a is a small GTPase that is involved in regulating cellular processes such as vesicular
295 transport and membrane trafficking (Guerra and Bucci, 2016). We hypothesized that the
296 observed reduction of ACE2 at the cell surface in *RAB7A* knock-out cells may be due to
297 impaired vesicular trafficking and accumulation of ACE2 in the cytoplasm. Immunofluorescence
298 in the A549^{ACE2} polyclonal cell line revealed that ACE2 is detected in most cells. We also found
299 that *RAB7A* knockout leads to an increased accumulation of ACE2 in the cytoplasm and in
300 vesicle-like hollow structures reminiscent of endo-lysosomes (**Figure 6C**). We detected these
301 accumulations in about 35% of *RAB7A* knockout cells with ACE2 staining, compared to the
302 control where ACE2 was primarily localized at the plasma membrane and small cytoplasmic
303 puncta (**Figure 6D**). Finally, we investigated which cellular compartments accumulate ACE2 in
304 *RAB7A* knockout cells. Co-immunofluorescence images showed that in *RAB7A* knockout cells
305 ACE2-containing vesicles often colocalize with EEA1, an early endosomal marker and less
306 frequently with LysoTracker, a lysosome marker (**Figure 6E**).

307
308 Since ACE2 was overexpressed in our A549^{ACE2} cells, we wondered whether Rab7a loss would
309 lead to similar cytoplasmic sequestration in cells with endogenous ACE2 expression. Flow
310 cytometry analysis in Caco-2 colon and Calu-3 lung cells revealed that *RAB7A* knock-out cells
311 have significantly reduced cell surface expression of ACE2 compared to cells transduced with a
312 non-targeting guide RNA (**Figure 6F, G**). We also found that *RAB7A* knock-out in Caco-2 cells

313 results in larger cytoplasmic ACE2 puncta compared to the control cells, further supporting
314 cytoplasmic accumulation of ACE2 in the absence of Rab7a (**Figure 6H, I**).

315

316 **Discussion**

317 Given the current COVID-19 global pandemic, there is an urgent need to better understand the
318 complex relationships between host and virus genetic dependencies. We report a genome-wide
319 loss-of-function screen in human lung cells that identified host genes required for SARS-CoV-2
320 viral infection. We selected and validated 30 genes that were ranked among the top 200 genes.
321 To support the ability of the screen to identify key dependencies, some of the well-known host
322 genes involved in SARS-CoV-2 Spike protein binding and entry such as the ACE2 receptor and
323 Cathepsin L were among the top-scoring genes (Hoffmann et al., 2020a). One of the validated
324 genes (*SIGMAR*) encodes the Sigma-1 receptor that was recently identified to be modulated by
325 drugs effective against SARS-CoV-2 *in vitro* (Gordon et al., 2020). Overall, the top-ranked
326 genes clustered within several protein complexes including vacuolar ATPases, Retromer and
327 endosome, Commander, ARP2/3, PI3K and others, highlighting both the critical importance of
328 multiple genes within each pathway to viral pathogenesis and the diversity of molecular
329 pathways involved in SARS-CoV-2 infection.

330

331 Using a “minipool” CRISPR library of perturbations targeting top-ranked genes from the
332 genome-scale CRISPR screen and single-cell transcriptomics, we identified a group of 6 genes
333 (*RAB7A*, *PIK3C3*, *NPC1*, *CCDC22*, *ATP6VIA*, and *ATP6API*) that had a similar transcriptional
334 signature — upregulation of the cholesterol synthesis pathway. By measuring the cholesterol
335 levels, we found that CRISPR-driven loss of those 6 genes result in increased cellular
336 cholesterol. Some of the 6 genes have previously been implicated in regulating low-density
337 lipoprotein (LDL) cholesterol. For example, depletion of Rab7a leads to LDL accumulation in
338 endosomes and NPC1 knockout cells show a reduction of cholesterol at the plasma membrane
339 and an accumulation in the late endosome/lysosome compartments (Chang et al., 2005; Girard et
340 al., 2014; Millard et al., 2000; Neufeld et al., 1996). We have recently, in an independent study,
341 identified that SARS-CoV-2 infection negatively downregulates the cholesterol synthesis
342 pathway and that viral infection can be counteracted by drug treatments that upregulate the same

343 pathway (Hoagland et al., 2020). It is possible that changes in lipid composition directly impacts
344 SARS-CoV-2 virion maturation and infectivity, as has been previously shown for Hepatitis C
345 and Influenza A (Aizaki et al., 2008; Bajimaya et al., 2017). In this study, we showed that
346 amlodipine, a calcium-channel antagonist, upregulates cholesterol levels and blocks SARS-CoV-
347 2 infection. In addition, recent clinical studies have suggested that patients taking amlodipine or
348 similar dihydropyridine calcium channel inhibitors have a reduced COVID-19 case fatality rate
349 (Solaimanzadeh, 2020; Zhang et al., 2020). An important future research direction will be to
350 further understand the relationship between cholesterol synthesis pathways and SARS-CoV-2.

351
352 Furthermore, we screened a panel of the top-ranked genes and identified that Rab7a regulates
353 cell surface expression of ACE2, likely by sequestering ACE2 in endosomal vesicles. Rab7a is
354 involved in vesicular trafficking and its depletion has been shown to sequester other cell
355 receptors in endosomes (Rush and Ceresa, 2013). Interestingly, *RAB7A* knock-out cell lines
356 showed both altered cholesterol biosynthesis and sequestration of ACE2 receptor. Also, previous
357 proteomics work showed that Rab7a has a strong interaction with viral protein nsp7 (Gordon et
358 al., 2020). However, there is no nsp7 in the incoming virion, implying a post-entry/post-
359 translational role for Rab7a. Thus, it is possible that loss of Rab7a blocks SARS-CoV-2
360 pathogenesis via multiple separate pathways, which is supported by the observation that it is the
361 top-performing gene in our arrayed validation.

362
363 While this study was under review, a few other groups released preprints with loss-of-function
364 CRISPR screens to identify host factors required for SARS-CoV-2 infection (Heaton et al., 2020;
365 Wei et al., 2020; Zhu et al., 2020). Notably, only two studies (ours and Zhu *et al.*) have
366 substantial overlap (11 and 14 genes among the top 36 genes from our MOI 0.3 and 0.01 screens,
367 respectively, as shown in **Figure S1D** and nearly all of the top-ranked gene categories shown in
368 **Figure 2A**). Given that these independent screens utilized different CRISPR libraries, the
369 corroboration by Zhu *et al.* provide further support for our conclusions. The overlap between
370 either of these studies with another screen performed using African green monkey cells (Wei *et*
371 *al.*) was limited to ACE2 and CTSL, two genes with established roles in viral entry. The
372 differences in the overlap among the top-ranked genes might be due to technical aspects (such as
373 different CRISPR libraries or variations in guide representation) or biological differences (such

374 as different cell types or different host species). An exciting avenue for future study would be to
375 investigate if SARS-CoV-2 perhaps utilizes multiple cell type-specific genetic circuits.

376
377 A key element in our study was harnessing genome-scale loss-of-function to develop more
378 refined therapeutic hypothesis. Our study suggests that PIK3C3 is a promising drug target: Four
379 out of the seven PIK3C3 inhibitors resulted in more than 100-fold reduction of SARS-CoV-2
380 viral load (SAR405, Compound-19, PIK-III, and Autophinib). Using a PIK3C3 polyclonal
381 knockout A549^{ACE2} cell line, we found that among the top four PIK3C3 inhibitors, SAR405 may
382 have some off-target effects (**Figure S4B**). Considering that our polyclonal knockout line likely
383 has some residual PIK3C3, future work will be required to test the inhibitor specificity using a
384 PIK3C3 monoclonal knockout cell line. Another drug that shows a substantial reduction in
385 SARS-CoV-2 viral load is tamoxifen. Tamoxifen is an FDA-approved drug given as prophylaxis
386 to patients at risk of breast cancer and works via modulation of the estrogen receptor. Tamoxifen
387 was included in our study as it targets protein kinase C as a secondary target (O'Brian et al.,
388 1985). This mechanism is further supported by the observation that A549 cells have undetectable
389 transcript levels of estrogen receptor 1 (Human Protein Atlas) (Uhlen et al., 2010). Considering
390 that tamoxifen is typically given to patients for years as a cancer therapy and prophylactic
391 (Marchant, 1976), it would be interesting to investigate if patients taking tamoxifen have a
392 reduced risk of SARS-CoV-2 infection and/or display less severe symptoms post-infection.

393
394 Finally, many approaches for therapeutic discovery have focused on large-scale screens of
395 compound libraries. Even when promising therapeutic candidates are identified, it can be
396 challenging to understand the mechanisms responsible for reducing viral pathogenesis. Our
397 forward-genetics approach allows us to first identify key host genes, which can then be targeted
398 through a diversity of methods such as small-molecule inhibitors, blocking antibodies or gene
399 knockdown. A key advantage of this approach is that the mechanism of action for any
400 therapeutic is well-established from the outset.

401
402 Taken together, our integrative study identifies essential host genes in SARS-CoV-2 viral
403 pathogenesis and, through a broad range of analytic and experimental approaches, validates their
404 central role in infection. We also identify potential mechanisms underlying top-ranked genes,

405 including cholesterol synthesis and endosomal function. In addition to guiding new therapeutic
406 targets to help end this pandemic, our study provides a framework for harnessing massively-
407 parallel genome editing to understand disease genetics and mechanisms.

408

409 **Limitations of Study**

410 Our study identified host factors required for SARS-CoV-2 infection in human A549 cells that
411 overexpress ACE2. Future work will be needed to explore limitations of our study. (1)
412 Considering the ACE2 overexpression in our screen, it will be interesting to screen human cells
413 expressing endogenous ACE2, which may potentially identify transcriptional regulators of
414 ACE2. (2) The A549 cell line used in our CRISPR screen is a lung adenocarcinoma cell line.
415 Given that various organs are affected by SARS-CoV-2, it will be helpful to understand whether
416 there are tissue-specific host factors. (3) While we show through multiple, distinct genetic
417 perturbations that upregulation of the cholesterol biosynthesis pathway and increase in cellular
418 cholesterol blocks SARS-CoV-2, the precise mechanisms of how changes in cholesterol disrupt
419 viral infection remain to be elucidated. (4) Recent genome-wide association studies have
420 uncovered human genetic variants associated with COVID-19 risk and severity. Since the
421 majority of such variants are in noncoding regions, integrative analysis of genome-wide CRISPR
422 screens may help pinpoint the causal genes through which these variants function.

Acknowledgements

We thank the entire Sanjana and tenOever laboratories for support and advice. We also thank the New York Genome Center for flow cytometry and confocal microscopy resources and T. Maniatis for advice. Z.D. is supported by an American Heart Association postdoctoral fellowship (grant no. 20POST35220040). Postdoctoral fellowship support for T.X.J. is provided by the NIH (grant no. R01AI123155). M.L. is supported by the Hope Funds for Cancer Research postdoctoral fellowship. T.L. is supported by NIH/NIMH (R01MH106842) and NIH/NHGRI (UM1HG008901). S.K. and T.L. are supported by NIH/NHLBI (R01HL142028). B.R.t. is supported by the Marc Haas Foundation, the National Institutes of Health, and DARPA's PREPARE Program (HR0011-20-2-0040). N.E.S. is supported by NYU and NYGC startup funds, NIH/NHGRI (DP2HG010099), NIH/NCI (R01CA218668), DARPA (D18AP00053), the Sidney Kimmel Foundation, and the Brain and Behavior Foundation.

Author Contributions

Z.D., T.X.J., B.R.t. and N.E.S. conceived the project. Z.D., T.X.J., B.R.t. and N.E.S. designed and performed the genome-scale CRISPR screen. N.E.S., Z.D., H-H.W. and E.G. analyzed the CRISPR screen. Z.D., T.X.J., D.A.H., M.L., L.L., B.R.t. and N.E.S. designed and performed the arrayed validation and inhibitor work. T.X.J. performed BSL3 work. O.D. and B.R.R. made the transgenic A549 cells. H-H.W., Z.D., E.P.M., P.S. and N.E.S. designed and performed the single-cell experiments. S.M., Z.D. and N.E.S. designed and performed the endosomal trafficking experiments. S.K. and T.L. assisted with genetic analyses. N.E.S., B.R.t., H.P., P.S. and T.L. supervised the work. Z.D., T.X.J., H-H.W., D.A.H., B.R.t. and N.E.S. wrote the manuscript with input from all the authors.

Declaration of Interests

N.E.S. is an advisor to Vertex. T.L. is a member of the scientific advisory boards of Goldfinch Bio and, with equity, Variant Bio. The New York Genome Center and New York University have applied for patents relating to the work in this article.

Main figure titles and legends

Figure 1. A genome-scale CRISPR loss-of-function screen to identify genes that prevent SARS-CoV-2 infection of human alveolar epithelial carcinoma cells.

A) Overview of the genome-scale loss-of-function screen for host factors in human A549^{ACE2} cells requires for SARS-CoV-2 infection. **B)** Immunofluorescence of SARS-CoV-2 nucleocapsid (N) protein and DAPI labeling of human A549^{ACE2} cells at 24 hours post-infection. **C)** Percent survival of human A549^{ACE2} cells transduced with the GeCKOv2 library with the indicated SARS-CoV-2 viral amount (MOI) at 6 days post-infection. **D)** Scatterplot of guide RNA read counts from A549^{ACE2} cells at 6 days post-infection with SARS-CoV-2 (MOI ~ 0.01) versus cells prior to infection. Read counts are normalized log₂ reads. **E)** Volcano plot of median fold-change of guide RNAs for each gene and log₁₀ Robust Rank Aggregation (RRA) *p*-values. All genes with |fold-change| > 4 and RRA *p* < 10⁻³ are labeled. **F)** Overlap of top 50 ranked genes between the MOI 0.01 and MOI 0.3 screen.

Figure 2. Top-ranked genes from the CRISPR screen are involved in key elements of the SARS-CoV-2 viral lifecycle.

Schematic of SARS-CoV-2 docking, entry, RNA genome release and transcription, and virion replication, assembly and release with top-ranked host genes from the CRISPR screen highlighted in red. All genes shown are ranked in the top 50 genes (top ~0.25% of library) in the low MOI CRISPR screen using Robust Rank Aggregation. Adapted from Du *et al.* (2009).

Figure 3. Enriched genes cluster into related pathways, are expressed broadly, interact directly with viral proteins, and are also involved in viral pathogenesis of pandemic flu and Zika virus.

A) Classification of genes shown in Figure 2 (top-ranked ~0.25% of the GeCKOv2 library) into specific complexes. **B)** Gene set enrichment analysis normalized enrichment scores for all significant (FDR *q* < 0.1) Gene Ontology (GO) biological processes. **C)** Expression of top-ranked genes (same as in A) across the indicated human tissues from GTEx v8. Gene expression color scale is transcripts per million (TPM). **D)** RRA fold-change for the low MOI CRISPR screen for the high-confidence protein-protein interaction with the maximum fold-change for each viral gene from the Gordon *et al.* mass spectrometry dataset (2020). **E)** Clustering of top-ranked GO biological processes for CRISPR screens for Zika virus ZIKV (Y. Li *et al.*, 2019), H1N1 pandemic avian influenza IAV (B. Li *et al.*, 2020), and SARS-CoV-2 (this study).

Figure 4. Arrayed validation of genome-scale SARS-CoV-2 screen and identification of druggable gene targets.

A) Representative immunofluorescence images of A549^{ACE2} knockout lines infected with SARS-CoV-2 at MOI of 0.1 and fixed 36 hours post-infection (hpi). SARS-CoV-2 N protein is shown in red and DAPI in blue. **B)** Quantification of SARS-CoV-2 infected A549^{ACE2} knockout lines immuno-stained with N protein as shown in A. Each gene was targeted with 3 different guide RNAs represented as diamond symbols ($n = 3$ biological replicates, error bars indicate s.e.m.). **C)** Correlation of \log_2 median fold change from the genome-scale CRISPR screen (low MOI) and percent of infected cells after individual (arrayed) gene perturbation shown in B. **D)** Druggable genes found in the Drug Gene Interaction database (DGIdb) among highly-ranked genes from the genome-scale CRISPR screen (ranked by RRA p -value). **E)** Quantitative PCR (qPCR) of SARS-CoV-2 viral load present in A549^{ACE2} cells pretreated for 2 hours with the indicated small molecule inhibitors at 10 μ M and then infected with SARS-CoV-2 at MOI of 0.1. The qPCR was performed at 36 hours post-infection (hpi). Red bars indicate inhibitors that yield a greater than 100-fold reduction in viral load. Bars with hatch marks indicate an unreliable viral load measurement due to a large reduction in cell viability (see panel F). Inhibitors were maintained at the same concentration throughout the experiment ($n = 6$ biological replicates, error bars indicate s.e.m.). **F)** Percent of A549^{ACE2} viable cells following inhibitor treatments at 10 μ M for 36 hpi determined using LIVE/DEAD stain and flow cytometry. Bars with red hatch marks indicate that inhibitor treatment had a large impact on viability ($< 90\%$ viability). Significance testing for panels B, E, and F was performed via a one-way ANOVA with false-discovery rate-corrected follow up tests; for clarity of presentation, all significance testing can be found in **Table S6**.

Figure 5. Single-cell transcriptomics (ECCITE-seq) identifies shared target gene signatures for lipid and cholesterol regulation.

A) Schematic of pooled CRISPR perturbations with Expanded CRISPR-compatible Cellular Indexing of Transcriptomes and Epitopes by sequencing (ECCITE-seq). Adapted from Mimitou *et al.* (2019). **B)** Single-cell mRNA expression heatmap showing the 100 most differentially upregulated genes (adjusted p -value < 0.01) for 200 randomized cells per selected target gene perturbation (for clarity, CCDC22, PIK3C3, RAB7, and TMEM165 perturbations are not shown). Labeled genes are either a top-ranked gene from the genome-wide CRISPR screens (*red*) or among the top 5 differentially-expressed genes for a gene perturbation (*black*). **C)** Heatmap summarizing Gene Set Enrichment Analysis results for genes upregulated in any of the indicated target gene-perturbed cells (all genes with p -value < 0.01 and with a limit to the 300 most differentially-expressed genes; all enriched pathways with adjusted p -value $< 10^{-13}$).

D) Cholesterol quantification in gene-perturbed cells (with the indicated guide RNAs), normalized to total protein.

Figure 6. RAB7A loss results in a reduced cell surface expression and an increased endosomal accumulation of ACE2.

A) Representative histograms of flow cytometry analysis to determine cell surface expression of ACE2 on A549 cell lines (A549 wild type (WT), A549^{ACE2}, and ACE2 with Cas9 and non-targeting (NT) or RAB7A-targeting guide RNAs). The dashed line indicates the gate between the ACE2 negative and positive cells. **B)** Fraction of ACE2+ cells (using gating shown in panel A). ACE2 expression level was normalized across all samples to the A549^{ACE2} cells transduced with non-targeting (NT) guides ($n = 2-3$ gRNA-transduced lines per gene, error bars are s.e.m.). **C)** Representative images of immunofluorescence staining of ACE2 on A549^{ACE2} transduced with a non-targeting (NT) or a RAB7A-targeting guide. In NT cells, ACE2 localizes at the cell membrane and in the cytoplasm, while in RAB7A-targeted cells, ACE2 shows a distinct pattern of localization to vesicles. **D)** Percent of cells with ACE2 accumulation in vesicles in NT and RAB7A-transduced A549^{ACE2} cells ($n = 2$ biological replicates, error bars are s.e.m.). **E)** Representative images of immunofluorescence co-stained for ACE2, EEA1 and LysoTracker in A549^{ACE2} cells with a CRISPR guide RNA targeting RAB7A. ACE2 shows a distinct colocalization with EEA1 (an early endosome marker) and a less frequent colocalization with LysoTracker (a lysosomal marker). **F)** Representative histograms of flow cytometry analysis to determine cell surface expression in Calu-3 cells. The dashed line indicates the gate between the ACE2 negative and positive cells. **G)** Fraction of ACE2+ cells (using gating shown in panel F). ACE2 expression level was normalized across all samples to the Calu-3 cells transduced with a non-targeting (NT) guide ($n = 3$ biological replicates, error bars are s.e.m.). **H)** Representative images of immunofluorescence staining of ACE2 on Caco-2 cells transduced with a NT or a RAB7A-targeting guide. In NT cells, ACE2 localizes at the cell membrane and in the cytoplasm, while in RAB7A-targeted cells, ACE2 shows a distinct pattern of localization to vesicles. **I)** Mean area of ACE2 foci in Caco-2 cells transduced with a NT or a RAB7A-targeting guide ($n = 4$ biological replicates, 80-105 cells per replicate were scored, error bars are s.e.m.). Significance testing for panels B and G was performed with a one-way ANOVA (B: $F = 9.8$, $p < 10^{-4}$; G: Calu-3: $F = 378$, $p < 10^{-4}$, Caco-2: $F = 222$, $p < 10^{-4}$) with false-discovery rate-corrected post-hoc tests. Significance testing for panels D and I was performed with an unpaired t -test. For all panels, * indicates $p \leq 0.05$, ** indicates $p \leq 0.01$, *** indicates $p \leq 0.001$, and **** indicates $p \leq 0.0001$.

Supplemental figure titles and legends

Supplementary Figure 1. Genome-wide loss-of-function CRISPR screen enriched gene identification, Related to Figure 1.

A) Scatterplot of guide RNA read counts from A549^{ACE2} cells at 3 days post-transduction with the GeCKOv2 library versus read counts from the library plasmid. Read counts are normalized log₂ reads. B) RRA *p*-value distribution for all genes in the GeCKOv2 library. C) Overlap of top-ranked (top 500) genes between 3 different analysis methods (RRA, RIGER, and SBR). 142 genes are found by all 3 methods. D) Comparison of top-ranked genes between different genome-scale CRISPR screens for SARS-CoV-2 infection.

Supplementary Figure 2. Gene set enrichment and overlap of top-ranked genes with other viral infections, Related to Figures 2 and 3.

A) – D) Four of the significant (FDR < 0.1) top-ranked GO biological process terms and the fold-change ranks of their genes in the SARS-CoV-2 low MOI CRISPR screen. E) Normalized gene ranks of the top 50 genes from the SARS-CoV-2 low MOI CRISPR screen and genome-scale CRISPR screens for Zika virus (ZIKV) and H1N1 avian influenza (IAV).

Supplementary Figure 3. Validation of top-ranked genes using CRISPR perturbations and RNA interference, Related to Figure 4.

A) Western blot analysis of RAB7A, CCDC22, ATP6V1A, and ACE2 after transduction of A549^{ACE2} with the indicated CRISPR guide RNA and selection with puromycin for 7 days. For validation, we designed 3 independent guide RNAs per gene (i.e. distinct guide RNAs from those in the GeCKOv2 library). Beta tubulin was used as loading control. B) Quantitative PCR (qPCR) of SARS-CoV-2 viral load present in A549^{ACE2} CRISPR-perturbed cells infected with SARS-CoV-2 at MOI of 0.1. The qPCR was performed on cells collected at the indicated time (hours) post-infection (hpi) (*n* = 6 biological replicates, error bars indicate s.e.m.). C) Western blot analysis of RAB7A and ACE2 after transduction of Huh7.5^{ACE2} with the indicated CRISPR guide RNA and selection with puromycin for 7 days. For validation, we designed 3 independent guide RNAs per gene (i.e. distinct guide RNAs from those in the GeCKOv2 library). Beta tubulin was used as loading control. D) qPCR of SARS-CoV-2 viral load present in Huh7.5^{ACE2} CRISPR-perturbed cells infected with SARS-CoV-2 at MOI of 0.1. The qPCR was performed on cells collected and fixed at 36 h.p.i (*n* = 3 guide RNAs with 6 biological replicates each, error bars indicate s.e.m.). E) Immunofluorescence quantification of SARS-CoV-2 N protein at 36 hours post-infection (hpi) at MOI 0.1 in A549^{ACE2} cells pretreated with siRNA pools for 48 hours (*n* = 3 technical replicates, error bars represent s.e.m., NT indicates non-targeting controls).

Supplementary Figure 4. Perturbations of enriched CRISPR screen genes with small-molecule inhibitors, Related to Figure 4.

A) Immunofluorescence quantification of SARS-CoV-2 N protein at 36 hpi (MOI 0.1) in A549^{ACE2} cells pretreated for 2 hours with 10 μ M of the indicated inhibitors ($n = 3$ biological replicates, error bars represent s.e.m.) **B)** Quantitative PCR (qPCR) of SARS-CoV-2 viral load present in A549^{ACE2} cells (CRISPR-perturbed with either non-targeting or PIK3C3-targeting guide RNAs) pretreated for 2 hours with the indicated PIK3C3 molecule inhibitors at 10 μ M and then infected with SARS-CoV-2 at MOI of 0.1. The qPCR was performed at 36 hours post-infection (hpi). Inhibitors were maintained at the same concentration throughout the experiment ($n = 6$ biological replicates, error bars indicate s.e.m.). **C)** Immunofluorescence quantification of SARS-CoV-2 N protein at 36 hpi (MOI 0.1) in A549^{ACE2} cells pretreated with a combination of indicated inhibitors at 10 μ M each for 2 hours ($n = 3$ biological replicates, error bars represent s.e.m.).

Supplementary Figure 5. ECCITE-seq identifies cholesterol gene signature shared across multiple top-ranked genes, Related to Figure 5.

A) Stacked violin plot of 11 genes shared in both ECCITE-seq experiments. Single-cells are grouped by unique guide RNA target gene label of cells with a single detected guide RNA. Target gene expression is highlighted in red. **B)** Heatmap of Gene Set Enrichment Analysis results for genes downregulated in any of the indicated target gene perturbed cells. **C)** Cholesterol (normalized by total protein) in A549^{ACE2} cells treated with amlodipine or vehicle (DMSO) for 24 hours. **D)** Quantitative PCR (qPCR) of SARS-CoV-2 viral load present in A549^{ACE2} cells treated with amlodipine or DMSO for 24 hours and then infected with SARS-CoV-2 at MOI of 0.1. The qPCR was performed on cells collected at the indicated time (hours) post-infection (hpi) ($n = 3$ biological replicates, error bars indicate s.e.m.). **E)** Plaque assays of SARS-CoV-2 viral load present in A549^{ACE2} cells treated with amlodipine or DMSO for 24 hours and then infected with SARS-CoV-2 using logarithmically diluted supernatants. ($n = 3$ biological replicates, error bars indicate s.e.m.). **F)** Number of reads mapping to the indicated portion of the viral genome in A549^{ACE2} cells treated with amlodipine or DMSO. A representative sample is shown for each treatment ($n = 3$ biological replicate sequencing libraries). **G)** Cell viability by Trypan Blue exclusion in A549^{ACE2} cells treated with amlodipine or DMSO for 24 hours ($n = 3$ biological replicates, error bars indicate s.e.m.). **H)** Distance matrix of RNA-sequencing from A549^{ACE2} cells treated with amlodipine or DMSO for 24 hours and then infected at MOI 0.1 or mock infection ($n = 3$ biological replicate sequencing libraries for each treatment-infection group). Read counts were processed with the DESeq2 regularized-log transform before computing distances. **I)** k -means clustering ($k = 3$) of the top 500 most variable

genes across all 4 conditions ($n = 3$ biological replicate sequencing libraries for each treatment-infection group). For each cluster, we label the most enriched pathway (lowest p -value) and, for the genes in that pathway, we label the top 5 most variable genes. No significantly enriched pathways were found for Cluster 3.

Supplementary Figure 6. Flow cytometry for cell surface ACE2 expression and protein analysis of RAB7A protein after CRISPR targeting, Related to Figure 6.

A) and **B)** Flow cytometry gating strategy to quantify cell surface expression of ACE2. **A)** Live cells were first gated by the forward and side scatter area, then doublets were excluded by gating with the forward scatter area and width. Viable cells were selected by gating on side scatter area and LIVE/DEAD violet. **B)** Gating strategy to determine ACE2⁺ cells. The gate was position such that <3% of A549 wild type and >85% of A549^{ACE2} cells were ACE2 positive. The same gating strategy was applied to all samples. **C)** Western blot on A549^{ACE2} cells perturbed with non-targeting (NT) or RAB7A-targeting guide RNAs and probed with a RAB7A antibody. GAPDH was used as loading control.

STAR Methods

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Neville Sanjana (neville@sanjanalab.org).

Materials Availability

The pHR-PGK-ACE2 vector generated in this study is available through Addgene (161612).

Data and Code Availability

CRISPR screen, single cell RNA-sequencing/ECCITE-seq and bulk RNA-sequencing datasets are available on the GEO repository with accession numbers GSE158298, GSE159519 and GSE159522, respectively. Additional Supplemental Items are available from Mendeley Data at <http://dx.doi.org/10.17632/km23bwyny6.1>

Experimental Models and Subject Details

Mammalian cell lines and culture conditions

Human alveolar basal epithelial carcinoma cells (A549, ATCC CCL-185), human hepatocellular carcinoma (Huh7.5, a kind gift from C. Rice), human colorectal carcinoma (Caco-2, ATCC HTB-37), lung adenocarcinoma (Calu-3, ATCC HTB-55), monkey kidney epithelial cells (Vero E6, ATCC CRL-1586) and human embryonic kidney cells HEK293FT (Thermo) were used in our study. HEK293FT, Huh7.5, and Caco-2 cells were maintained at 37°C and 5% CO₂ in D10 media, which consists of DMEM (Caisson Labs) with 10% Serum Plus II Medium Supplement (Sigma-Aldrich). Calu-3 and Vero E6 cells were maintained in EMEM (ATCC) media with 10% Serum Plus II Medium Supplement (Sigma-Aldrich).

Viral strains

SARS-related coronavirus 2 (SARS-CoV-2), isolate USA-WA1/2020 (NR-52281), used in the study (Blanco-Melo et al., 2020; Daniloski et al., 2020). SARS-CoV-2 was grown in Vero E6 cells in DMEM supplemented with 2% FBS, 4.5 g/L D-glucose, 4 mM L-glutamine, 10 mM non-essential amino acids, 1 mM sodium pyruvate and 10 mM HEPES. Plaque assays were used to determine infectious titers of SARS-CoV-2 by infection of Vero E6 cells in Minimum Essential Media supplemented with 2% FBS, 4 mM L-glutamine, 0.2% BSA, 10 mM HEPES and 0.12% NaHCO₃ and 0.7% agar.

Method Details

Generation of A549^{ACE2} line

To generate ACE2 expressing cells, the human ACE2 coding sequence was amplified and cloned into the BamHI site of the lentiviral vector pHR-PGK (Addgene 79120). We generated lentiviral particles (as described below) and transduced 5×10^4 A549 cells plated in a 12-well plate in the presence of polybrene (8 μ g/ml). We confirmed hACE2 expression by western blot analysis (Thermo MA5-32307).

Lentiviral production, transduction and SARS-CoV-2 infection

The Human GeCKOv2 A and B libraries (Addgene 1000000048) were used for genome-scale CRISPR knockout screens (Sanjana et al., 2014). We mixed equal amount of the A and B library plasmids to target each gene with 6 guide RNAs. Briefly, 225cm² flasks of 80% confluent HEK293FT cells (Thermo) were transfected with 25 μ g GeCKOv2 plasmid, 14 μ g pMD2.G and 20 μ g psPAX2 mixed in 2.5 mL OptiMEM (Thermo) and 175 μ L Polyethylenimine (1 mg/ml) (Polysciences). After 6 hours, media was changed to D10 media with 1 % bovine serum albumin (Sigma) added to improve virus stability. After 60 hours, viral supernatants were harvested and centrifuged at 3,000 rpm at 4 °C for 10 min to pellet cell debris debris and filtered using 45 μ m PVDF filters (CellTreat). The supernatant was then ultracentrifuged for 2 hours at 100,000g (Sorvall Lynx 6000) and the pellet resuspended overnight at 4 °C in PBS with 1% BSA.

Following lentiviral titration, 330 million A549^{ACE2} expressing cells were transduced at MOI of 0.5. Cells were selected with 2 μ g/uL puromycin (Thermo) for 12 days to ensure proper selection. Throughout the experiment the representation was monitored such that each guide RNA is represented by at least 1000 cells (~125 M cells).

We infected GeCKOv2-transduced cells (~1000x representation) with SARS-CoV-2 isolate USA-WA1/2020 (NR-52281) at either MOI of 0.01 (low MOI) and 0.3 (high MOI). Surviving cells were collected on day 6 post-infection for genomic DNA isolation.

Genomic DNA isolation, guide RNA amplification and quantification

We used a two-step PCR protocol (PCR1 and PCR2) to amplify the guide RNA cassette for Illumina sequencing from genomic DNA (gDNA). The gDNA was extracted from CRISPR screen cells using the following protocol (Chen et al., 2015): Per 100 million cells, 12 mL of NK Lysis Buffer (50 mM Tris, 50 mM EDTA, 1% SDS, pH 8) were used for cell lysis. Once cells were resuspended, 60 μ L of 20 mg/ml Proteinase K (Qiagen) was added and the sample was incubated at 55°C overnight. The next day, 60 μ L

of 20 µg/mL RNase A (Qiagen) was added, mixed, and samples were incubated at 37°C for 30 min. Then 4 mL of pre-chilled 7.5 M ammonium acetate was added, samples were vortexed and spun at 4,000g for 10 min. The supernatant was placed in a new tube, mixed well with 12 mL isopropanol and spun at 4,000g for 10 min. DNA pellets were washed with 12 mL of 70% ethanol, spun, dried and pellets were resuspended with 0.2x TE buffer (Sigma).

For the first PCR reaction, we used all gDNA available for each sample. We performed the PCR1 using Taq-B polymerase (Enzymatics) and used multiple reactions where each reaction contained up to 10 µg gDNA per 100 µL PCR reaction. PCR1 products for each sample were pooled and used for amplification with barcoded PCR2 primers. For each sample, we performed 12 PCR2 reactions (using 5 µL of the pooled PCR1 product per PCR2 reaction) with Q5 polymerase (NEB). PCR2 products were pooled and then normalized within each biological sample before combining uniquely-barcoded separate biological samples. The pooled product was then gel-purified from a 2% E-gel EX (Life Technologies) using the QiaQuick gel extraction kit (Qiagen). The purified, pooled library was then quantified with TapeStation 4200 (Agilent Technologies). PCR products were run on a 2% agarose gel and the correct size band was extracted. Sequencing was performed on the NextSeq 550 instrument using the HighOutput Mode v2 with 75 bp paired-end reads (Illumina). All PCR1 and PCR2 primer sequences, including full barcodes, are listed on the GeCKO website (<http://genome-engineering.org/gecko/>).

Sequencing reads were demultiplexed upon sequencing based on Illumina i7 barcodes present in PCR2 reverse primers using Illumina BaseSpace. We performed adaptor trimming by treating the hU6 promoter sequence as a 5' adapter, using `cutadapt v1.13 [-e 0.2 -o 5 -m 20 -g TCTTGTGGAAAGGACGAAACACCG]`. Processed guide RNA sequences were aligned to the GeCKOv2 reference allowing for up to 1 mismatch using `bowtie v1.1.2 [-a --best --strata -v 1 -norc]` with alignment rates of 81% to 86%.

Computational analyses of genome-scale CRISPR screens

Guide RNA counts were processed using the MaGeCK pipeline with an output of RRA *p*-values and gene ranks (Chen et al., 2018). We separately ranked genes using the RIGER (weighted-sum) and second-best rank methods (Chen et al., 2015; Luo et al., 2008). Gene Set Enrichment Analyses were performed using the `fgsea` package with Gene Ontology for biological processes (`c5.bp.v7.1.symbols`) (Korotkevich et al., 2019). GTEx v8 tissue specific enrichment was performed using the Multi Gene Query function available on the GTEx website: <https://www.gtexportal.org/home/multiGeneQueryPage> (accessed August 1st, 2020) (Aguet et al., 2019). GO enrichments for SARS-CoV-2, ZIKV, and IAV CRISPR screens were performed using `GOrilla` to find all significant enrichments (FDR *p*-value < 10-

3) (Eden et al., 2009). FDR p -values were \log_{10} -transformed and normalized to create the heatmap shown in **Figure 3E**.

Generation of gene-perturbed human cell lines and SARS-CoV-2 infection

For validation by CRISPR knock-out (using 3 independent guide RNAs per gene), we selected genes that were highly-ranked (either in the low or high MOI screen or both) or genes that had not previously been implicated in viral pathogenesis or immune signaling. For each gene of interest, 3 guide RNAs were designed using GUIDES software and were subsequently cloned into an all-in-one vector with an optimized Cas9 scaffold (pCC_01, Addgene 139086) containing Cas9, a guide RNA cassette and a puromycin resistant cassette (Legut et al., 2020; Meier et al., 2017). Following sequence confirmation by Sanger sequencing, lentivirus was produced individually for each plasmid and the target cells were transduced in presence of polybrene (10 $\mu\text{g}/\text{mL}$, Santa Cruz). Gene-perturbed A549^{ACE2}, Huh7.5^{ACE2}, Calu-3 and Caco-2 cells were selected for at least 10 days with 2 $\mu\text{g}/\text{mL}$ (A549^{ACE2}, Calu-3), 3 $\mu\text{g}/\text{mL}$ (Caco-2) or 5 $\mu\text{g}/\text{mL}$ (Huh7.5^{ACE2}) puromycin (Thermo). To determine the SARS-CoV-2 infection susceptibility of gene-perturbed lines, 10,000 cells were plated per well of 96-well plates. After 24 hours, the cells were infected with SARS-CoV-2 at MOI of 0.1. At 36 hours post-infection, the cells were either fixed and processed for immunofluorescence or cellular RNA was harvested for qRT-PCR analyses. For the multistep growth curves, cells were infected at an MOI of 0.1, and total RNA was harvested from infected cells at indicated times. RNA was processed for qRT-PCR as described above. All infections with SARS-CoV-2 were performed with at least 3 biological replicates.

Immunofluorescence of nucleocapsid (N) protein

Cells were fixed with 5% formaldehyde and immunostained for nucleocapsid (N) protein and visualized with a secondary antibody labeled with AlexaFluor-568 (Thermo). SARS-CoV-2 nucleocapsid (N) antibody (clone 1C7C7) was obtained from the Center for Therapeutic Antibody Discovery at the Icahn School of Medicine at Mount Sinai. Nuclei were stained with DAPI. Full wells were imaged and quantified for SARS-CoV-2 infected cells using a Celigo imaging cytometer (Nexcelom Biosciences). All infections with SARS-CoV-2 were performed with 3 biological replicates. Representative images from the top gene knockout hits were acquired using the EVOS M5000 Imaging System (Thermo).

Identification of druggable genes and drug treatments

To identify druggable genes among the top CRISPR screen hits, we cross-referenced highly-ranked genes from the RRA analysis with the data table containing drugs and their gene targets was obtained from Drug Gene Interaction database (DGIdb, retrieved on June 3, 2020) as well as manual literature search

(Cotto et al., 2018). Inhibitors that target the genes of interest (and remdesivir) was obtained from SelleckChem and MedChemExpress. The catalog number and vendor information is available in the Key Resource section. Amlodipine was obtained from Sigma (A5605).

To test drug efficacy in reducing SARS-CoV-2 infection, 10,000 A549^{ACE2} cells were seeded per well of a 96-well plate. Cells were treated with inhibitors at 10 μ M for two hours before infection and inhibitors were maintained throughout the course of infection. Cells were infected with SARS-CoV-2 at MOI of 0.1 for 36 hours and the cells were collected for analysis via qRT-PCR or processed for immunofluorescence (N protein and quantified by Celigo). For Figure 4E, the remdesivir data was collected in an independent experiment.

Cell viability assays

Cell viability following drug treatments was performed the same way as described above. Thirty-six hours post drug treatment the cells were collected and stained with LIVE-DEAD Violet (Thermo). Cell acquisition was performed using a Sony SH800S cell sorter with a 100 μ m sorting chip. We used the following gating strategy: 1) We excluded the cell debris based on the forward and reverse scatter; 2) Doublets were excluded by plotting FSC height vs FCS area, 3) Dead cells were quantified using live-dead violet stain. For all samples, we recorded at least 5,000 cells that pass the gating criteria described above. Flow cytometry analyses were performed using FlowJo v10. Cell viability for A549-ACE2 treated with DMSO or 10 μ M amlodipine was performed 24 hours post treatment using Trypan Blue (Thermo Fisher) and automated cell counting (Nexcelom AutoT4).

Quantitative reverse-transcription PCR (qRT-PCR) of viral RNA

RNA was extracted from cells grown in 96-well plates by using the RNeasy 96 Kit (Qiagen) per the manufacturer's instructions. RNA was reverse-transcribed and PCR amplified using SYBR Fast One-step Universal qRT-PCR Kit (KAPA/Roche). For amlodipine experiments, RNA was extracted using TRIzol Reagent (Invitrogen) and purified with the Direct-zol RNA Miniprep kit (Zymo Research) as per the manufacturer's instructions (including the optional DNase I treatment). For amlodipine experiments, the RNA was reverse-transcribed using SuperScript Reverse Transcriptase II (Thermo Fisher) and oligo(dT) primers. The cDNA was diluted 1:20 before qRT-PCR was performed using SYBR Fast qPCR (KAPA/Roche). SARS-CoV-2 replication was assessed by using primers specific to the N mRNA (Forward 5'-CTCTTGATAGATCTGTTCTCTAAACGAAC-3'; Reverse 5'-GGTCCACCAAACGTAATGCG-3'). SARS-CoV-2 N mRNA levels were normalized to beta tubulin (Forward 5'-GCCTGGACCACAAGTTTGAC-3; Reverse 5'-TGAAATTCTGGGAGCATGAC-3'). Reactions were ran and analyzed on a Lightcycler 480 II Instrument (Roche). Relative quantification was calculated by

comparing the cycle threshold (C_t) values using $\Delta\Delta C_t$. C_t values of subgenomic nucleocapsid transcript were normalized to C_t values of alpha-tubulin, and those values were normalized to viral transcript levels of DMSO-treated or non-targeting guide RNA-transduced control samples, and displayed as fold-reduction in viral transcript. Significance was determined using a one-way ANOVA with post-hoc tests or a two-tailed unpaired Student's t -test, as appropriate.

siRNA transfections and SARS-CoV-2 infection

All of the siRNAs were ordered from Dharmacon and their catalog number can be found in the Key Resource Table. To knockdown individual genes, 10,000 A549^{ACE2} cells were seeded in 96-well plates and transfected with siRNAs using Lipofectamine RNAiMAX (Thermo) following the manufacturer's protocol. Forty-eight hours later, the cells were infected with SARS-CoV-2 at an MOI of 0.1 for 36 hours. Cells were fixed with 5% formaldehyde, stained with nucleocapsid protein (clone 1C7C7, ISMMS), and visualized with AlexaFluor-568 conjugated secondary antibody (Thermo). Nuclei were stained with DAPI, and full wells were imaged with a Celigo imaging cytometer (Nexcelom Biosciences).

Minipool CRISPR library and lentiviral production

To generate the minipool, we combined equimolar amounts the same guide RNA vectors (cloned in pCC1) that we used for arrayed validation (see above *Generation of gene-perturbed A549^{ACE2} lines and SARS-CoV-2 infection*). In total, we combined 3 guide RNAs for each gene (either, 30 or 11 genes) plus 9 non-targeting guide RNAs. The non-targeting guide RNA plasmids were added at half the molar ratio of the other plasmids. Lentivirus was produced via transfection of the minipool with appropriate packaging plasmids (psPAX2: Addgene 12260; pMD2.G: Addgene 12259) using polyethylenimine (PEI) reagent in HEK293FT. The target A549^{ACE2} cells were transduced in presence of polybrene (10 $\mu\text{g}/\text{mL}$, Santa Cruz) (ECCITE-Seq experiment 1) or without (ECCITE-seq experiment 2) at very low MOI to ensure entry of a single virus per cell. The survival was below 5% after 2 days of selection with 2 $\mu\text{g}/\text{mL}$ puromycin (Thermo) for both ECCITE-seq experiments. On day 10 post-transduction, the cells were collected and processed for ECCITE-seq.

Expanded CRISPR-compatible Cellular Indexing of Transcriptomes and Epitopes (ECCITE-seq)

To perform cell-hashing and ECCITE-seq (Mimitou et al., 2019; Stoeckius et al., 2018), we separated cells into several sub-pools and, for each pool, we resuspended 1 million cells in 100 μl staining buffer (2% BSA, 0.01% Tween in PBS). We then added 10 μl Human TruStain FcX Fc Receptor Blocking Solution (BioLegend) and incubated on ice for 10 minutes. We then added hashing antibodies (BioLegend), incubated on ice for 30 minutes, and washed cells 3 times with staining buffer. In total, we

ran four lanes of a Chromium Single Cell Immune Profiling Solution v1.0 5' kit (10x Genomics) targeting recovery of 12,000 cells per lane (superloading) in experiment 1 and 20,000 cells in experiment 2. Gene expression (mRNA), hashtags (Hashtag-derived oligos, HTOs) and guide RNA (Guide-derived oligos, GDOs) libraries were constructed (Smibert et al., 2019). Each replicate was sequenced on two NextSeq550 75-cycle high-output runs (Illumina). Sequencing reads from the mRNA library were mapped to the hg38 reference genome (Ensembl v97) using Cellranger (v3.0.1, 10x Genomics). To generate count matrices for HTO and GDO libraries, the CITE-seq-count package was used (<https://github.com/Hoohm/CITE-seq-Count> v1.4.2). Count matrices were then used as input into the Seurat R package (v3.2) for downstream analyses (Stuart et al., 2019).

ECCITE-seq data analysis

Processing of initial ECCITE-seq experiment with 30 target genes: Cells with low quality metrics, high mitochondrial gene content ($> 17.5\%$) and low number of genes detected (≤ 1800) were removed. The median number of detected genes was 3309. RNA counts were log-normalized and HTO counts were normalized using the centered log-ratio transformation approach, with `margin=2` (normalizing across cells). To identify cell doublets and assign experimental conditions to cells, we used the `HTOseqDemux` function (Stoeckius et al., 2018). `HTOseqDemux`-defined cell doublets and negatives were removed from any downstream analyses.

Cellular guide RNA identity for cells in the ECCITE-seq pool was assigned based on GDO unique molecular identifier (UMI) counts. We considered a guide RNA detected with ≥ 16 UMI counts (the median UMI counts per guide RNA was 503 counts, 93% of cells with ≥ 1 sgRNA). We observed that 64% of the cells had >1 detected guide RNA (29% 1 guide RNA, 27% 2 guide RNAs, 17% 3 guide RNAs). To maximize cell recovery, we retained cells with 1 - 3 guide RNAs per cell ($n = 10,265$). We collapsed the guide RNA information down to target gene level (3 guide RNA per target gene) in the following way: For all cells (with 1-3 guide RNAs), we assigned a unique target gene if all detected guide RNAs targeted the same gene, or if the additional guide RNA detected was a non-targeting (NT) guide RNA (4,715 cells: 4,013 cells with exactly 1 sgRNA and 702 cells with collapsed target gene assignment). All remaining cells assigned 2 or 3 separate target genes were discarded.

Some of the top-ranked genes from the CRISPR screens may be endpoint genes and thus may have subtle phenotypes difficult to detect using single-cell transcriptomics. To address this issue, we used the initial ECCITE-seq experiment ($n = 4,715$ cells) and grouped cells by target gene. To identify target gene perturbations that lead to systemic transcriptomic changes, we determined the number of differentially expressed genes relative to cells with NT guide RNAs. We found that under stringent filtering conditions

(non-adjusted p -value $< 10^{-5}$ or adjusted p -value < 0.2), only 11 target genes showed more than 5 differentially expressed genes. The second ECCITE-seq experiment was conducted using guide RNAs solely for these.

Joint processing of both ECCITE-seq experiments: Cells with low quality metrics, high mitochondrial gene content ($> 15\%$) and low or high number of genes detected (≤ 2000 and > 5000) were removed. The median number of detected genes was 3365, with a median of 13,532 UMIs per cell. HTO counts were normalized using the centered log-ratio transformation approach, with `margin=2` (normalizing across cells). To identify cell doublets and assign experimental conditions to cells, we used the `HTOseqDemux` function (Stoeckius et al., 2018). `HTOseqDemux`-defined cell doublets and negatives were removed from any downstream analyses. Guide RNA counts were normalized similar to HTO UMI counts. Cell-wise guide RNA assignment was achieved using the `MultiSeqDemux` function (McGinnis et al., 2019). We only retained cells assigned with a single HTO and a single guide RNA (ECCITE-seq experiment 1: $n = 1,824$ cells, ECCITE-seq experiment 2: $n = 17,029$ cells). Cells from ECCITE-Seq experiment 1 and experiment 2 were harmonized by log-normalizing, followed by the identification of the 2000 most variable features with “vst” selection method. Next, we used `FindIntegrationAnchors` and `IntegrateData` with the first 40 principle components.

In the harmonized data set ($n = 18,853$ cells), we used `FindMarkers` to find differentially expressed genes between non-targeting cells and cells that belonged to a targeted gene class. We used up to 100 of the most upregulated genes with adjusted p -value < 0.01 and minimal \log_eFC threshold > 0.1 filter as input into the heatmap in **Figure 5B**. The same `FindMarkers` analysis (but using the non-adjusted p -value < 0.01 with a maximum of the 300 most significant DEGs) was used as input into the `EnrichR` package to run pathway analysis using the human WikiPathways database (v. 2019) (Kuleshov et al., 2016; Slenter et al., 2018). **Figure 5C** shows $-\log_{10}$ transformed p -values for the union of all enriched pathways with an adjusted p -value < 0.01 across the 11 target genes with perturbation signatures.

Plaque assay

Infectious titer in supernatant of SARS-CoV-2 infections (from A549^{ACE2}) was determined by plaque assay in Vero E6 cells. Cells were seeded in 12-well plates before infection with logarithmically diluted supernatants in a total inoculum of 100 μ l, rocking plates every 10 minutes for 1 hour. Then an overlay of Minimum Essential Media with 0.12% NaHCO_3 , 4 mM L-glutamine, 0.2% BSA, 10 mM HEPES and 0.7% OXOID agar (Thermo) was aliquoted into wells and plates were incubated for 48 hours before fixing with 5% formaldehyde. After 24 hours, plaques were stained with crystal violet (1% crystal violet in 20% EtOH) for 15 minutes before washing wells and counting plaques.

Bulk RNA-sequencing

RNA-sequencing libraries were constructed from 1 μ g of RNA per sample using the TruSeq Stranded mRNA Library Prep Kit (Illumina) according to the manufacturer's instructions. Sequencing libraries were sequenced using an Illumina NextSeq 500 platform and fastq files were generated using `bcl2fastq` (Illumina). Alignment to the SARS-CoV-2 genome (GenBank: MN985325.1) were performed using `bowtie` (Langmead et al., 2009). Reads were aligned to hg19 using STAR aligner in the Basespace RNA-Seq Alignment application (Illumina) and processed using DESeq2 (Love et al., 2014). Sample distances were calculated using read counts from the regularized log transform in DESeq2. K-means clustering was performed using the `kmeans` function in R/RStudio. Enriched Reactome pathways were identified using the STRING tool (Szklarczyk et al., 2019).

Measurement of cellular cholesterol

Cellular cholesterol was measured using the Cholesterol-Glo assay (Promega). For each cell line (CRISPR-perturbed or drug-treated), we counted 100,000 cells, washed with 1xPBS twice and resuspended in 100 μ L cell lysis solution for 30 min at 37°C. Following the incubation, 10 μ L of cell lysate was used to measure the cholesterol levels in 96 well plates in a final reaction volume of 100 μ L per well (without esterase). Luminescence was measured after incubating the plate for 1 hour at room temperature in the dark. Using the same lysis sample, the protein concentration was determined using the BCA assay (Thermo). The cholesterol levels were normalized to the respective protein concentration of each sample. A549^{ACE} cells were incubated with 10 μ M amlodipine (Sigma) or DMSO for 24 hours. Cholesterol levels were then determined as described above.

Flow cytometry of ACE2 cell surface expression

Cells were harvested, counted and about 100,000 cells were washed with Dulbecco's phosphate-buffered saline (PBS, Caisson Labs) and then stained with LIVE/DEAD Violet stain (Thermo, 34864). Following a wash with 1x PBS, all subsequent washes and antibody dilutions were performed using 1x PBS supplemented with 2% FBS. Wild-type or Cas9-perturbed A549^{ACE2}, Caco-2 and Calu-3 cells were stained for 30 min on ice with 0.25 μ g of anti-ACE2 antibody (R&D Systems, AF933) in ~50 μ L residual volume. Following two washes with 1x PBS with 2% FBS, samples were stained on ice for 20 min with 7.5 μ L of anti-goat-APC secondary antibody (R&D Systems, F0108). Cell acquisition and sorting was performed using a Sony SH800S cell sorter with a 100 μ m sorting chip. We used the following gating strategy: 1) We excluded the cell debris based on the forward and reverse scatter; 2) Doublets were excluded by plotting FSC height vs FCS area, 3) Dead cells were excluded by live-dead stain. We recorded at least 5,000 cells for A549^{ACE2} and at least 3,000 cells for Calu-3 and Caco-2 that pass the

gating criteria described above. Gates to determine ACE2-APC⁺ cells were set based on control A549 wild type cells or only secondary antibody stained Calu-3 and Caco-2, where the percent of ACE2 positive cells was set as <5% (background level). Flow cytometry analyses were performed using FlowJo v10.

Immunofluorescence of ACE2 and endo/lysosomal markers

A549^{ACE2} Cas9-transduced (specific gene or non-targeting) cell lines were seeded on poly-D-lysine-coated coverslips (Electron Microscopy Sciences) 24 hours before they were fixed with 4% formaldehyde (Sigma) diluted in 1x PBS (Caisson Labs) (Daniloski et al., 2020). Following 3 washes with 1x PBS, cells were blocked with 1x PBS with 2% BSA for 30 min at room temperature. Cells were stained with mouse anti-2A antibody diluted at 1:250 (clone 3H4, Millipore Sigma, MABS2005) to recognize ACE2-2A, and rabbit anti-EEA1 diluted at 1:100 (Thermo, MA5-14794). To detect ACE2 in Caco-2 cells, goat anti-ACE2 (R&D Systems, AF933) was used at 1.25 µg/mL. Cells were incubated with primary antibodies for 3 hours at room temperature in a moisturized chamber. Coverslips were then washed 5 times with 1x PBS with 2% BSA and then incubated with Alexa-conjugated secondary antibodies (Jackson ImmunoResearch) diluted at 1:1000 for 45 min at room temperature. Coverslips were then washed 5 times with 1x PBS with 2% BSA, and DAPI (Sigma) was added in the fourth wash at a concentration of 0.1 µg/mL. To stain the lysosomes, LysoTracker DeepRed (Thermo) was diluted to 70 nM in culture media and was added to actively growing cells on coverslips and were incubated for 1 hour at 37°C. Following incubation with LysoTracker, cells were processed for immunofluorescence as described above. Confocal images were acquired on a Zeiss LSM 780 with a 20x/0.8 Plan-Apochromat objective or a 63x/1.4 Plan-Apochromat objective (Zeiss). Epifluorescence images were acquired using the same system; both fitted with a Zeiss Axiocam 506 mono (Zeiss). Images were processed using Zen Black 2012 (Zeiss) and FIJI 2.1.0; Java 1.8.0-202 (Schindelin et al., 2012). Linear histogram adjustments were applied uniformly within experiments for clarity of presentation.

ACE2 localization was determined on images taken on a Zeiss LSM 780 with a 20x/0.8 Plan-Apochromat objective using Zen Black (Zeiss). The A549^{ACE2} polyclonal cell line had both ACE2 positive and negative cells. Only ACE2 positive cells were then manually scored if they had a distinct accumulation of cytoplasmic vesicles-like hollow structures, compared to the non-targeting (NT) control. In the NT control, the ACE2 staining was seen on the plasma membrane and diffuse in the cytoplasm. The scoring was performed on 2 biological replicates (that is, two separate non-targeting controls and two separate Rab7A knockout lines engineered using 4 separate Cas9 guide RNAs). We scored ~430 cells per replicate. To quantify the size of ACE2 foci, confocal microscopy images were thresholded using FIJI's

local-threshold (Phansalkar radius = 15) and then a watershed transform was applied. Foci were counted and measured using FIJI's `Analyze Particles` function.

Western blots

A549^{ACE2} or Huh7.5^{ACE2} cells were collected, washed with 1x PBS and lysed with TNE buffer (10 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Nonidet P-40) in presence of protease inhibitor cocktail (Bimake B14001) for 1 hour on ice. Cells lysates were spun for 10 min at 10,000 g, and protein concentration was determined with the BCA assay (Thermo). Equal amounts of cell lysates (20µg) were denatured in Tris-Glycine SDS Sample buffer (Thermo LC2676), and loaded on a Novex 4 - 20% Tris-Glycine gel (Thermo XP04205BOX). PageRuler pre-stained protein ladder (Thermo 26616) was used to determine the protein size. The gel was run in 1x Tris-Glycine-SDS buffer (IBI Scientific IBI01160) for about 120 min at 120V. Proteins were transferred on a nitrocellulose membrane (BioRad 1620112) in presence of prechilled 1x Tris-Glycine transfer buffer (FisherSci LC3675) supplemented with 20% methanol for 100 min at 100V. Immunoblots were blocked with 5% skim milk dissolved in 1x PBS with 1% Tween-20 (PBST) and incubated overnight at 4°C separately with the following primary antibodies: rabbit anti-RAB7A (0.1µg/mL, NovusBio, NBP1-87174), rabbit anti-GAPDH (0.1 µg/mL, Cell Signaling, 2118S), rabbit anti-CCDC22 (0.34 µg/mL, Proteintech, 16636-1-AP), rabbit anti-ATP6V1A (0.46 µg/mL, Proteintech, 17115-1-AP), rabbit anti-ACE2 (0.5 µg/mL, Invitrogen, MA5-32307), and mouse anti-beta tubulin (0.5 µg/mL, Invitrogen, 32-2600). Following the primary antibody, the blots were incubated with IRDye 680RD donkey anti-rabbit (0.2 µg/mL, LI-COR 926-68073) or with IRDye 800CW donkey anti-mouse (0.2 µg/mL, LI-COR 926-32212). The blots were imaged using Odyssey CLx (LI-COR).

Quantification and Statistical Analysis

Statistical analysis

Statistical parameters and details are reported in each figure legend. Generally, experiments were repeated with at least three biological replicates. Each plot includes points for individual biological replicates and mean \pm s.e.m. error bars unless otherwise specified. In general, biological replicates consist of replicate viral transductions (CRISPR), replicate RNA transfections (siRNA), or replicate drug treatments (small molecule inhibitors) and are specified in each figure legend. For imaging experiments, the number of cells scored per biological replicate is given in the figure legend.

When assessing significance of genetic perturbations and/or inhibitor treatments, we employed first a one-way or two-way ANOVA to assess significance and, when significant, post-hoc comparisons were performed using the false-discovery rate method of Benjamini-Krieger-Yekutieli to correct for multiple comparisons. Before applying ANOVA, we first verified the equality of group variances using the Brown-Forsythe test. Details for all ANOVA and post-hoc multiple comparisons are reported in **Table S6**; *t*-tests are reported in the respective figure legends.

Throughout the manuscript, we use r to denote the Pearson correlation and r_s to denote the Spearman correlation; all reported correlations were significant using the `cor.test` function in R. For the BioID interactome analysis, we derived an empirical p value by resampling from a null distribution ($n = 10,000$ gene shuffling iterations). Statistical analyses were performed in GraphPad Prism 8 and RStudio 1.2.5019.

Supplementary Tables

- Table S1.** SARS-CoV-2 pooled CRISPR screen ranked gene lists for low and high MOI screens, Related to Figures 1 and S1.
- Table S2.** Gene set enrichment analysis for SARS-CoV-2 pooled CRISPR screen; affinity proteomics enriched CRISPR screen genes for each viral gene; Gene Ontology (GO) analyses of SARS-CoV-2, ZIKV and IAV pooled CRISPR screens, Related to Figures 2, 3 and S2.
- Table S3.** Arrayed validation CRISPR guide RNAs and short-interfering RNAs, Related to Figures 4, S3 and S4 and STAR Methods.
- Table S4.** DGIdb druggable gene details for top-ranked CRISPR screen genes and FDA approval status of tested inhibitors, Related to Figures 4 and S4.
- Table S5.** ECCITE-seq reagents and differentially-expressed genes, Related to Figures 5 and S5 and STAR Methods.
- Table S6.** Significance testing results, Related to Figures 4, 5, S3 and S4.

References

- Aguet, F., Barbeira, A.N., Bonazzola, R., Brown, A., Castel, S.E., Jo, B., Kasela, S., Kim-Hellmuth, S., Liang, Y., Oliva, M., et al. (2019). The GTEx Consortium atlas of genetic regulatory effects across human tissues. *BioRxiv* 787903.
- Aizaki, H., Morikawa, K., Fukasawa, M., Hara, H., Inoue, Y., Tani, H., Saito, K., Nishijima, M., Hanada, K., Matsuura, Y., et al. (2008). Critical role of virion-associated cholesterol and sphingolipid in hepatitis C virus infection. *J. Virol.* 82, 5715–5724.
- Bajimaya, S., Frankl, T., Hayashi, T., and Takimoto, T. (2017). Cholesterol is required for stability and infectivity of influenza A and respiratory syncytial viruses. *Virology* 510, 234–241.
- Banerjee, S., and Kane, P.M. (2020). Regulation of V-ATPase Activity and Organelle pH by Phosphatidylinositol Phosphate Lipids. *Front Cell Dev Biol* 8, 510.
- Beigel, J.H., Tomashek, K.M., Dodd, L.E., Mehta, A.K., Zingman, B.S., Kalil, A.C., Hohmann, E., Chu, H.Y., Luetkemeyer, A., Kline, S., et al. (2020). Remdesivir for the Treatment of Covid-19 - Preliminary Report. *N. Engl. J. Med.*
- Blanco-Melo, D., Nilsson-Payant, B.E., Liu, W.-C., Uhl, S., Hoagland, D., Møller, R., Jordan, T.X., Oishi, K., Panis, M., Sachs, D., et al. (2020). Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. *Cell* 181, 1036-1045.e9.
- Bouhaddou, M., Memon, D., Meyer, B., White, K.M., Rezelj, V.V., Correa Marrero, M., Polacco, B.J., Melnyk, J.E., Ulferts, S., Kaake, R.M., et al. (2020). The Global Phosphorylation Landscape of SARS-CoV-2 Infection. *Cell* 182, 685-712.e19.
- Chang, T.-Y., Reid, P.C., Sugii, S., Ohgami, N., Cruz, J.C., and Chang, C.C.Y. (2005). Niemann-Pick type C disease and intracellular cholesterol trafficking. *J. Biol. Chem.* 280, 20917–20920.
- Chen, C.-H., Xiao, T., Xu, H., Jiang, P., Meyer, C.A., Li, W., Brown, M., and Liu, X.S. (2018). Improved design and analysis of CRISPR knockout screens. *Bioinformatics* 34, 4095–4101.
- Chen, S., Sanjana, N.E., Zheng, K., Shalem, O., Lee, K., Shi, X., Scott, D.A., Song, J., Pan, J.Q., Weissleder, R., et al. (2015). Genome-wide CRISPR screen in a mouse model of tumor growth and metastasis. *Cell* 160, 1246–1260.
- Cotto, K.C., Wagner, A.H., Feng, Y.-Y., Kiwala, S., Coffman, A.C., Spies, G., Wollam, A., Spies, N.C., Griffith, O.L., and Griffith, M. (2018). DGIdb 3.0: a redesign and expansion of the drug-gene interaction database. *Nucleic Acids Res.* 46, D1068–D1073.
- Daniloski, Z., Jordan, T.X., Ilmain, J.K., Guo, X., Bhabha, G., tenOever, B.R., and Sanjana, N.E. (2020). The Spike D614G mutation increases SARS-CoV-2 infection of multiple human cell types. *BioRxiv* 2020.06.14.151357.
- Du, L., He, Y., Zhou, Y., Liu, S., Zheng, B.-J., and Jiang, S. (2009). The spike protein of SARS-CoV--a target for vaccine and therapeutic development. *Nat Rev Microbiol* 7, 226–236.
- Eden, E., Navon, R., Steinfeld, I., Lipson, D., and Yakhini, Z. (2009). GOrilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* 10, 48.
- Erb, M.A., Scott, T.G., Li, B.E., Xie, H., Paulk, J., Seo, H.-S., Souza, A., Roberts, J.M., Dastjerdi, S., Buckley, D.L., et al. (2017). Transcription control by the ENL YEATS domain in acute leukaemia. *Nature* 543, 270–274.
- Funk, C.D., Laferrière, C., and Ardakani, A. (2020). A Snapshot of the Global Race for Vaccines Targeting SARS-CoV-2 and the COVID-19 Pandemic. *Front Pharmacol* 11, 937.
- Gardner, L. (2020). Johns Hopkins Coronavirus Resource Center.
- Gates, B. (2020). Responding to Covid-19 - A Once-in-a-Century Pandemic? *N. Engl. J. Med.* 382, 1677–1679.
- Girard, E., Chmiest, D., Fournier, N., Johannes, L., Paul, J.-L., Védie, B., and Lamaze, C. (2014). Rab7 is functionally required for selective cargo sorting at the early endosome. *Traffic* 15, 309–326.

- Gordon, D.E., Jang, G.M., Bouhaddou, M., Xu, J., Obernier, K., White, K.M., O'Meara, M.J., Rezelj, V.V., Guo, J.Z., Swaney, D.L., et al. (2020). A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature* 583, 459–468.
- Guerra, F., and Bucci, C. (2016). Multiple Roles of the Small GTPase Rab7. *Cells* 5.
- Heaton, B.E., Trimarco, J.D., Hamele, C.E., Harding, A.T., Tata, A., Zhu, X., Tata, P.R., Smith, C.M., and Heaton, N.S. (2020). SRSF protein kinases 1 and 2 are essential host factors for human coronaviruses including SARS-CoV-2. *BioRxiv* 2020.08.14.251207.
- Hoagland, D.A., Clarke, D.J.B., Møller, R., Han, Y., Yang, L., Wojciechowicz, M.L., Lachmann, A., Oguntuyo, K.Y., Stevens, C., Lee, B., et al. (2020). Modulating the transcriptional landscape of SARS-CoV-2 as an effective method for developing antiviral compounds. *BioRxiv* 2020.07.12.199687.
- Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.-H., Nitsche, A., et al. (2020a). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 181, 271-280.e8.
- Hoffmann, M., Kleine-Weber, H., and Pöhlmann, S. (2020b). A Multibasic Cleavage Site in the Spike Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. *Mol. Cell* 78, 779-784.e5.
- Jain, I.H., Zazzeron, L., Goli, R., Alexa, K., Schatzman-Bone, S., Dhillon, H., Goldberger, O., Peng, J., Shalem, O., Sanjana, N.E., et al. (2016). Hypoxia as a therapy for mitochondrial disease. *Science* 352, 54–61.
- Kim, D., Lee, J.-Y., Yang, J.-S., Kim, J.W., Kim, V.N., and Chang, H. (2020). The Architecture of SARS-CoV-2 Transcriptome. *Cell*.
- Kolde, R., Laur, S., Adler, P., and Vilo, J. (2012). Robust rank aggregation for gene list integration and meta-analysis. *Bioinformatics* 28, 573–580.
- Korotkevich, G., Sukhov, V., and Sergushichev, A. (2019). Fast gene set enrichment analysis. *BioRxiv* 060012.
- Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A., et al. (2016). Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res.* 44, W90-97.
- Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10, R25.
- Legut, M., Daniloski, Z., Xue, X., McKenzie, D., Guo, X., Wessels, H.-H., and Sanjana, N.E. (2020). High-throughput screens of PAM-flexible Cas9 variants for gene knock-out and transcriptional modulation. *Cell Reports* 2020.01.22.916064.
- Lek, A., Zhang, Y., Woodman, K.G., Huang, S., DeSimone, A.M., Cohen, J., Ho, V., Conner, J., Mead, L., Kodani, A., et al. (2020). Applying genome-wide CRISPR-Cas9 screens for therapeutic discovery in facioscapulohumeral muscular dystrophy. *Sci Transl Med* 12.
- Li, B., Clohisey, S.M., Chia, B.S., Wang, B., Cui, A., Eisenhaure, T., Schweitzer, L.D., Hoover, P., Parkinson, N.J., Nachshon, A., et al. (2020). Genome-wide CRISPR screen identifies host dependency factors for influenza A virus infection. *Nat Commun* 11, 164.
- Li, Y., Muffat, J., Omer Javed, A., Keys, H.R., Lungjangwa, T., Bosch, I., Khan, M., Virgilio, M.C., Gehrke, L., Sabatini, D.M., et al. (2019). Genome-wide CRISPR screen for Zika virus resistance in human neural cells. *Proc. Natl. Acad. Sci. U.S.A.* 116, 9527–9532.
- Liu, Y., Gayle, A.A., Wilder-Smith, A., and Rocklöv, J. (2020). The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J Travel Med* 27.
- Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.

- Luo, B., Cheung, H.W., Subramanian, A., Sharifnia, T., Okamoto, M., Yang, X., Hinkle, G., Boehm, J.S., Beroukhi, R., Weir, B.A., et al. (2008). Highly parallel identification of essential genes in cancer cells. *Proc. Natl. Acad. Sci. U.S.A.* *105*, 20380–20385.
- Mallam, A.L., and Marcotte, E.M. (2017). Systems-wide Studies Uncover Commander, a Multiprotein Complex Essential to Human Development. *Cell Syst* *4*, 483–494.
- Marchant, J. (1976). Letter: Prophylactic chemotherapy for breast cancer. *Br Med J* *1*, 1589.
- McGinnis, C.S., Patterson, D.M., Winkler, J., Conrad, D.N., Hein, M.Y., Srivastava, V., Hu, J.L., Murrow, L.M., Weissman, J.S., Werb, Z., et al. (2019). MULTI-seq: sample multiplexing for single-cell RNA sequencing using lipid-tagged indices. *Nat. Methods* *16*, 619–626.
- McNally, K.E., and Cullen, P.J. (2018). Endosomal Retrieval of Cargo: Retromer Is Not Alone. *Trends Cell Biol.* *28*, 807–822.
- Meier, J.A., Zhang, F., and Sanjana, N.E. (2017). GUIDES: sgRNA design for loss-of-function screens. *Nat. Methods* *14*, 831–832.
- Millard, E.E., Srivastava, K., Traub, L.M., Schaffer, J.E., and Ory, D.S. (2000). Niemann-pick type C1 (NPC1) overexpression alters cellular cholesterol homeostasis. *J. Biol. Chem.* *275*, 38445–38451.
- Mimitou, E.P., Cheng, A., Montalbano, A., Hao, S., Stoeckius, M., Legut, M., Roush, T., Herrera, A., Papalexi, E., Ouyang, Z., et al. (2019). Multiplexed detection of proteins, transcriptomes, clonotypes and CRISPR perturbations in single cells. *Nat. Methods* *16*, 409–412.
- Mori, S., Ito, H., and Yamamoto, K. (1988). Effects of calcium antagonists on low density lipoprotein metabolism in human arterial smooth muscle cells. *Tohoku J. Exp. Med.* *154*, 329–333.
- Neufeld, E.B., Cooney, A.M., Pitha, J., Dawidowicz, E.A., Dwyer, N.K., Pentchev, P.G., and Blanchette-Mackie, E.J. (1996). Intracellular trafficking of cholesterol monitored with a cyclodextrin. *J. Biol. Chem.* *271*, 21604–21613.
- O'Brian, C.A., Liskamp, R.M., Solomon, D.H., and Weinstein, I.B. (1985). Inhibition of protein kinase C by tamoxifen. *Cancer Res.* *45*, 2462–2465.
- Patel, S.J., Sanjana, N.E., Kishton, R.J., Eidizadeh, A., Vodnala, S.K., Cam, M., Gartner, J.J., Jia, L., Steinberg, S.M., Yamamoto, T.N., et al. (2017). Identification of essential genes for cancer immunotherapy. *Nature* *548*, 537–542.
- Ranganathan, S., Harmony, J.A., and Jackson, R.J. (1982). Effect of Ca²⁺ blocking agents on the metabolism of low density lipoproteins in human skin fibroblasts. *Biochem. Biophys. Res. Commun.* *107*, 217–224.
- Riva, L., Yuan, S., Yin, X., Martin-Sancho, L., Matsunaga, N., Pache, L., Burgstaller-Muehlbacher, S., De Jesus, P.D., Teriete, P., Hull, M.V., et al. (2020). Discovery of SARS-CoV-2 antiviral drugs through large-scale compound repurposing. *Nature*.
- Rush, J.S., and Ceresa, B.P. (2013). RAB7 and TSG101 are required for the constitutive recycling of unliganded EGFRs via distinct mechanisms. *Mol. Cell. Endocrinol.* *381*, 188–197.
- Samavarchi-Tehrani, P., Abdouni, H., Knight, J.D.R., Astori, A., Samson, R., Lin, Z.-Y., Kim, D.-K., Knapp, J.J., St-Germain, J., Go, C.D., et al. (2020). A SARS-CoV-2 – host proximity interactome. *BioRxiv* 2020.09.03.282103.
- Sanjana, N.E., Shalem, O., and Zhang, F. (2014). Improved vectors and genome-wide libraries for CRISPR screening. *Nat. Methods* *11*, 783–784.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods* *9*, 676–682.
- Shalem, O., Sanjana, N.E., Hartenian, E., Shi, X., Scott, D.A., Mikkelsen, T.S., Heckl, D., Ebert, B.L., Root, D.E., Duench, J.G., et al. (2014). Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* *343*, 84–87.

- Slenter, D.N., Kutmon, M., Hanspers, K., Riutta, A., Windsor, J., Nunes, N., Mélius, J., Cirillo, E., Coort, S.L., Digles, D., et al. (2018). WikiPathways: a multifaceted pathway database bridging metabolomics to other omics research. *Nucleic Acids Res.* *46*, D661–D667.
- Smibert, P., Mimitou, E., and Stoeckius, M. (2019). ECCITE-seq. Protocol Exchange.
- Solaimanzadeh, I. (2020). Nifedipine and Amlodipine Are Associated With Improved Mortality and Decreased Risk for Intubation and Mechanical Ventilation in Elderly Patients Hospitalized for COVID-19. *Cureus* *12*, e8069.
- Stoeckius, M., Zheng, S., Houck-Loomis, B., Hao, S., Yeung, B.Z., Mauck, W.M., Smibert, P., and Satija, R. (2018). Cell Hashing with barcoded antibodies enables multiplexing and doublet detection for single cell genomics. *Genome Biol.* *19*, 224.
- Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. *Cell* *177*, 1888–1902.e21.
- Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A.* *102*, 15545–15550.
- Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva, N.T., Morris, J.H., Bork, P., et al. (2019). STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* *47*, D607–D613.
- Uhlen, M., Oksvold, P., Fagerberg, L., Lundberg, E., Jonasson, K., Forsberg, M., Zwahlen, M., Kampf, C., Wester, K., Hober, S., et al. (2010). Towards a knowledge-based Human Protein Atlas. *Nat. Biotechnol.* *28*, 1248–1250.
- Verschueren, E., Von Dollen, J., Cimermancic, P., Gulbahce, N., Sali, A., and Krogan, N.J. (2015). Scoring Large-Scale Affinity Purification Mass Spectrometry Datasets with MiST. *Curr Protoc Bioinformatics* *49*, 8.19.1–8.19.16.
- Wei, J., Alfajaro, M.M., Hanna, R.E., DeWeirdt, P.C., Strine, M.S., Lu-Culligan, W.J., Zhang, S.-M., Graziano, V.R., Schmitz, C.O., Chen, J.S., et al. (2020). Genome-wide CRISPR screen reveals host genes that regulate SARS-CoV-2 infection. *BioRxiv* 2020.06.16.155101.
- Zhang, L., Sun, Y., Zeng, H.-L., Peng, Y., Jiang, X., Shang, W.-J., Wu, Y., Li, S., Zhang, Y.-L., Yang, L., et al. (2020). Calcium channel blocker amlodipine besylate is associated with reduced case fatality rate of COVID-19 patients with hypertension. *MedRxiv* 2020.04.08.20047134.
- Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., Si, H.-R., Zhu, Y., Li, B., Huang, C.-L., et al. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* *579*, 270–273.
- Zhu, Y., Feng, F., Hu, G., Wang, Y., Yu, Y., Zhu, Y., Xu, W., Cai, X., Sun, Z., Han, W., et al. (2020). The S1/S2 boundary of SARS-CoV-2 spike protein modulates cell entry pathways and transmission. *BioRxiv* 2020.08.25.266775.



Highlights (85 chars, including spaces, max per bullet)

- Genome-wide CRISPR knockout screen identifies host factors for SARS-CoV-2 infection
- Top-ranked genes include vacuolar ATPases, Retromer, Commander and Arp2/3 complex
- Validation using CRISPR knockout, RNA interference and small molecule inhibitors
- Reduced infection via increased cholesterol biosynthesis and sequestration of ACE2

In Brief


To identify potential therapeutic targets for SARS-CoV-2, Daniloski et al. conduct a genome-wide CRISPR screen in human lung epithelial cells. They identify genes and pathways required for SARS-CoV-2 infection, including the vacuolar ATPase proton pump, Retromer, and Commander complexes. Using single-cell transcriptomics, they identify upregulation of cholesterol biosynthesis as a common mechanism underlying viral resistance, in addition to ACE2 sequestration.

Identify genes required for SARS-CoV-2 infection




Validate top-ranked genes

CRISPR knock-out RNA interference Small molecule inhibitors



Mechanisms blocking SARS-CoV-2






Figure 1




Figure 2




Figure 3






Figure 4


A



B



C



D




Figure 5




Figure 6